



Effect of chlorfluazuron and pyriproxyfen on the antennal morphology, pheromone production and response of surviving adults of *Tribolium castaneum* treated at the LC₅₀ level during the pupal stage



Reda F.A. Bakr^{a, c, *}, Olfat M. El-Monairy^b, Nehad M. El-Barky^b, Nancy M.B. El-Shourbagy^b

^a Entomology Department, Faculty of Science, Ain Shams University, Cairo, Egypt

^b Entomology Department, Faculty of Science, Benha University, Al-kalyobya, Egypt

^c Biology Department, Faculty of Science, King Khalid University, Abha, Saudi Arabia

ARTICLE INFO

Article history:
Accepted 19 August 2013

Keywords:
Chlorfluazuron
Pyriproxyfen
Tribolium castaneum
Pheromone production
Antennal sensilla

ABSTRACT

Initial experiments were carried out to determine the dosage mortality response of the rust-red flour beetle *Tribolium castaneum* to two insect growth regulators, the chitin synthesis inhibitor chlorfluazuron and the juvenile hormone analogue pyriproxyfen. The response was measured as the proportion of adults developing from individuals treated during the pupal stage. For males, at the LC₅₀ level, chlorfluazuron was more toxic than pyriproxyfen with LC₅₀ values of 10.6 and 12.6 ppm respectively. For females, pyriproxyfen was more potent than chlorfluazuron with LC₅₀ values of 7.1 and 8.3 ppm.

Subsequent experiments were carried out using adults that had survived after treatment at the LC₅₀ level during the pupal stage. Both responses to pheromone and the production of pheromone, by adults of both sexes that had been treated during the pupal stages with pyriproxyfen, were significantly more affected than those treated with chlorfluazuron.

Both treatments caused abnormalities in the antennae of adults of both sexes.

Published by Elsevier Ltd.

1. Introduction

Insect pests of stored grains cause high economic losses in many countries of subtropical and tropical regions. The rust-red flour beetle, *Tribolium castaneum* (Herbst), is one of the most serious pests of flour and other cereal products in Egypt and other countries. Treatments to control insect pests in such stored products must be active against the pests, and safe to humans and environmentally-friendly organisms. One promising way to fulfill this need is through the use of insect growth regulators (IGRs).

The term IGR was introduced to describe a new class of bio-rational compounds. IGRs are divided into three main groups; juvenoids, which mainly affect larval metamorphosis by mimicking juvenile hormone; ecdysteroids, which affect molting, and chitin synthesis inhibitors (CSIs), which interfere with cuticle formation (Post et al., 1974).

Chlorfluazuron is a novel chitin synthesis inhibitor that belongs to benzoylphenylureas (BPU) group and acts as an anti-molting agent and inhibits biosynthesis of chitin of an important constituent in insect cuticle. The cuticle loses elasticity and there is abnormal endocuticular deposition and abortive molting (Dhadialla et al., 2005). It is also known as an active larvicide with a broad spectrum of activity on various insects, including Lepidoptera, Coleoptera, Homoptera, Hymenoptera, and Orthoptera (Bakr et al., 2005).

Pyriproxyfen is a new juvenile hormone analogue (JHA). It acts as an anti-JH which artificially enhances JH levels preventing insect development to the adult stage. It competes with JH in binding to the JH receptors or to the JH-carrier of proteins, and injuring the corpora allata cells or interfering with JH biosynthesis (Leighton et al., 1981). It is an effective pesticide against Hymenoptera, Dicotyledon, and Heteroptera (Mojaver and Bandani, 2010).

Pheromones provide a major mode of intraspecific communication in insects and act to elicit a specific behavioral or developmental response from other individuals of the same species (Nordlund, 1981).

The objective of this study was to clarify the possibilities of using IGRs and sex pheromone in pest control.

* Corresponding author. Entomology Department, Faculty of Science, Ain Shams University, Cairo, Egypt.

E-mail address: redabakr55@gmail.com (R.F.A. Bakr).

Table 1
Toxicity of the tested IGRs against pupal stage of *T. castaneum*.

IGRs	Chlorfluazuron		Pyriproxyfen	
	Male	Female	Male	Female
LC ₅₀ ppm	10.6	8.3	12.6	7.1
Slope	$(0.73 \pm 9.2) \times 10^{-3}$	$(0.72 \pm 8.73) \times 10^{-3}$	$(1.02 \pm 1.45) \times 10^{-3}$	$(0.9 \pm 1.07) \times 10^{-2}$
Chi square	3.8	4.7	7.2	8.1
Degree of freedom	3	3	3	3
Probabilities	0.23	0.17	0.05	-1.78×10^{-2}
Toxicity index at LC ₅₀	100	85.54	84.13	100

2. Materials and methods

2.1. Insect colony

A laboratory colony of the red flour beetle, *T. castaneum*, was maintained for many generations under constant conditions of temperature at 30 °C and 70% R.H. in the Department of Entomology, Benha University. The rearing medium was wheat flour mixed by weight with brewer's yeast (95: 5 w:w).

2.2. Treatments

Chlorfluazuron was prepared by aqueous dilution of an emulsifiable concentrate containing 50 g/L active constituent (Atabron) and pyriproxyfen from an emulsifiable concentrate containing 100 g/L active constituent (Admiral). Aqueous dilutions of both compounds were used to produce concentrations of 0.1, 0.5, 1, 5 and 10 ppm.

Pupae of uniform age were obtained using the sieve of 495 µm which separates pupae from adults. Pupae were segregated into males and females and were dipped for 10 s then were transferred into suitable media. Four replicates (25 pupae for each) were run for each concentration. In addition; a corresponding untreated control group was used. The pupae were examined after eight days. Percentage mortality was calculated on the base of the number of adult emerged in relation to the number of pupae per petri dish. Uncompleted emerging adults were counted as dead. To determine the effect of LC₅₀ of chlorfluazuron and pyriproxyfen against *T. castaneum*, subsequent experiments were carried out on adults treated at these dose levels during the pupal stage. All tests were performed at constant conditions of temperature at 30 °C and 70% r.h.

2.3. Pheromone production and response

Studies were carried out using a vial type of olfactometer similar to that used by Burkholder (1970). It consisted of a glass vial (15 × 1.5 cm), which had a rubber plug with a movable glass rod. The latter had a broad inner end at which a small piece of masking tape was fixed. The insect tested for pheromone production was held by the masking tape, while the one tested for response was placed on the bottom of the vial. The distance between the two insects was 4 cm. There were ten replicate vials set up in this way, each vial containing two individuals (a male and a female). The tested males and females were at the age range of 8–10 days old. Pheromone extraction was prepared by placing 30 treated beetles of the same sex and of known age (8–10 days) in one mL of hexane. The latter was confined in a screw-top glass vial of 5 mL in capacity. The tops of the vial caps were foil-lined to avoid solvent loss or contamination. The beetles were held in solvents for 24 h in a refrigerator, after which they were removed. Extracts were stored in a deep freezer at 20 °C until used. A vial containing 30 untreated

beetles was held under the same conditions and served as a control. The concentration of the female extracts was (0.3 µ) female equivalents (FE) per 10 µ solvent, according to Hussien (1982).

2.4. Scanning electron microscope studies on adult antennae

The fine structure and distribution of various types of antennal sensilla at the age range of 8–10 day old females and males resulting from treated pupae were compared with those of untreated individuals by using scanning electron microscopy (SEM). This was done using high vacuum mode at the Regional Center of Mycology and Biotechnology, Cairo, Egypt.

2.5. Statistical analysis

The results obtained were evaluated using one-way analysis of variance "ANOVA" (Snedecor, 1971) using Pro. Lab. Version 7.5. The statistical program was set at the 1% level of significance ($P < 0.01$). The data were subjected to Probit Analysis (Finney, 1971) to calculate LC₅₀ values.

3. Results

3.1. Toxicity of the tested insect growth regulators against the pupal stage

Chlorfluazuron exhibited higher toxic action than pyriproxyfen against male pupae of *T. castaneum*, giving LC₅₀ values of 10.6 and 12.6 ppm, respectively (Table 1). On the other hand, pyriproxyfen was more potent than chlorfluazuron against female pupae of *T. castaneum*, recording the lowest LC₅₀ value of 7.1 ppm.

Forming a toxicity index, when the male pupal stage treated with chlorfluazuron was considered as the standard rather than pyriproxyfen, the potency of pyriproxyfen was 84.31% lower than the standard. On the other hand, when the female pupal stage treated with pyriproxyfen was considered as the standard rather than chlorfluazuron, the potency of chlorfluazuron was 85.54% of the standard.

3.2. Effect of LC₅₀ of the tested IGRs on responsiveness and production of pheromones in male and female adults

Figures 1 and 2 illustrate that the responsiveness and production of pheromone in treated groups with pyriproxyfen were significantly higher than those treated with chlorfluazuron.

3.2.1. Male response behavior to female

For chlorfluazuron, when treated males were tested against treated females, the level of response was 60% compared to the control response of 82%. For pyriproxyfen, the responses were 52% and 84% respectively. The response behavior in the first group consisted of a sequence of increasing levels of excitation. The first

Download English Version:

<https://daneshyari.com/en/article/4517144>

Download Persian Version:

<https://daneshyari.com/article/4517144>

[Daneshyari.com](https://daneshyari.com)