



Original research article

Comparisons of blood biochemical parameters, digestive enzyme activities and volatile fatty acid profile between Meishan and Yorkshire piglets



Shouqing Ma, Meilei Ma, Chunlong Mu, Kaifan Yu, Weiyun Zhu*

Jiangsu Key Laboratory of Gastrointestinal Nutrition and Animal Health, Laboratory of Gastrointestinal Microbiology, College of Animal Science, Nanjing Agricultural University, Nanjing 210095, China

ARTICLE INFO

Article history:

Received 9 November 2015

Accepted 1 December 2015

Available online 12 December 2015

Keywords:

Blood biochemical parameters

Digestive enzyme activity

Volatile fatty acid

Meishan piglets

Yorkshire piglets

ABSTRACT

This study was conducted to compare physiological characteristics between Meishan and Yorkshire piglets in their early lives. Six healthy purebred Meishan sows and Yorkshire sows with close farrowing dates were used in this research. The piglets sucked their respective sow's milk for 14 days, then they were slaughtered to collect samples of blood, pancreas, contents of stomach, jejunum, cecum, colon as well as feces for analysis of blood biochemical parameters, digestive enzymes, and volatile fatty acid (VFA). The results showed that Yorkshire piglets had higher concentrations of high-density lipoprotein cholesterol (HDL-C) and total cholesterol (TC) ($P < 0.05$). Gastric lipase activity was higher in Meishan piglets but Yorkshire piglets had higher lactase activity ($P < 0.05$). The total VFA together with acetate and propionate in cecum and colon were higher in Meishan piglets than in Yorkshire piglets ($P < 0.05$), but acetate in jejunum and ratio of acetate to propionate in colon were lower in Meishan piglets than in Yorkshire piglets ($P < 0.05$). In conclusion, in early suckling period, significant differences exist in host metabolism and intestinal microbial metabolism between Meishan and Yorkshire piglets.

© 2015, Chinese Association of Animal Science and Veterinary Medicine. Production and hosting by Elsevier B.V. on behalf of KeAi Communications Co., Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Different pig breeds, especially obese and lean breeds, have been proved to be different in physiological and biochemical characteristics. He et al. (2012) analyzed the blood metabolites of obese Ningxiang pigs (a Chinese indigenous breed) and lean Duroc × Landrace × Yorkshire crossbred pigs after weaning using nuclear magnetic resonance-based metabolomics. The result indicated that significant differences existed in lipids synthesis, lipids oxidation, energy metabolism and amino acids metabolism between the two type pigs. Our previous study on the fecal microbial

community of Meishan and Yorkshire piglets showed that Meishan piglets had a stronger ability in volatile fatty acid (VFA) production, and Yang et al. (2014) found that more bacteria belonged to the Firmicutes phyla. Both studies suggest that there are discrepancies in metabolic characteristics between obese and lean piglets after weaning.

To date, only a few studies have focused on physiological characteristics in suckling period without solid feed intake. Kelly et al. (1991) showed that genetically identical piglets fed maternal milk from different breed sows displayed differences in lactase activity, villus height and crypt depth. It was also reported that genetically different piglets fed the same sow milk presented little variance in body weight gain in the late period (Rzasa et al., 2002). These results indicated that maternal milk and genetics affected piglet development in the suckling period. But there is no study regarding the differences on the physiological and biochemical characteristics between obese and lean piglets in early suckling period to our knowledge.

Based on these studies, we hypothesized that the early physiological and biochemical characteristics of obese Meishan (a Chinese indigenous breed) piglets and lean Yorkshire piglets were different.

* Corresponding author.

E-mail address: zhuweiyun@njau.edu.cn (W. Zhu).

Peer review under responsibility of Chinese Association of Animal Science and Veterinary Medicine.



Production and Hosting by Elsevier on behalf of KeAi

Thus, blood biochemical parameters, digestive enzyme activities and VFA concentrations of the two breeds were determined during suckling period.

2. Materials and methods

2.1. Animal management and experimental design

Six healthy purebred Meishan sows and Yorkshire sows with close farrowing date were recruited in this research. All of the sows had 3 to 4 times of accouchement. The sows were given the same type and amount of diets based on corn and soybean. The contents of corn, soybean, fish meal, wheat bran and premix were 63, 24, 3, 6 and 4%, respectively. The nutrient levels of crud protein, ether extracts and crude fiber were 20.67, 3.22 and 3.21%, respectively. The sows were reared individually in one barn with nursing pens, the conditions were identical. After delivery, all piglets were reared on nursing pens until the end of the experiment.

On day 14 postnatal, 1 piglet was randomly selected from each litter (a total of 12 piglets), weighed, and then slaughtered. Before slaughter, blood samples were collected from the vena cava and centrifuged at $1,400 \times g$ for 10 min at 4°C . Serum samples were collected and stored at -20°C for future analysis.

2.2. Digesta and feces collection

Digesta samples and feces were collected from stomach, anterior jejunum, cecum and colon, and then stored at -80°C for digestive enzymes and VFA analysis.

2.3. Blood biochemical parameters

Total protein (TP), albumin (ALB), globulin (GLO), glucose (GLU), urea, total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C) of serum were determined by automatic biochemistry analyzer (AU-400, Olympus, Japan) with kits according to manufacturer's instruction.

2.4. Digestive enzyme activities

Contents of stomach, jejunum and pancreas were prepared for digestive enzymes determination by mixing 1 g contents with 3 mL deionized water. Through centrifugation ($2,500 \times g$ for 10 min at 4°C), supernatant fluids were obtained to determine activities of amylase, lipase and protease by using colorimetric method according to the instruction of the manufacture (Nanjing Jiancheng Bioengineering Institute, Nanjing, China).

2.5. Volatile fatty acid concentrations

The determination of VFA concentrations was described as previous study (Qin, 1982). Briefly, 0.3-g fecal samples were put in tubes, then 1 mL deionized water was added to each tube. After centrifugation ($9,000 \times g$ for 10 min at 4°C), 500 μL supernatant fluids were obtained and 0.1 mL 25% (wt/vol) metaphosphoric acid were added to them. The mixtures were injected into gas chromatograph (GC-14B, Shimadzu, Japan) for VFA determination. Parameter settings of the instrument were as follows, temperatures of capillary pipette column, vaporization and detection were 130, 180, and 180°C , respectively. The pressures of nitrogen (the carrier gas), hydrogen and oxygen were 60, 50 and 50 kPa, respectively.

2.6. Statistical analysis

All the experimental data are shown as means \pm SEM. Data were analyzed using SPSS 17.0 software (SPSS Inc., Chicago, IL). Significant level was determined by *T*-test. We considered $P < 0.05$ as statistical significance.

3. Results

3.1. Comparisons of blood biochemical parameters between Meishan and Yorkshire piglets

Blood biochemical parameters measured in the current experiment are presented in Table 1. The concentrations of HDL-C and TC were significantly lower in Meishan piglets than in Yorkshire piglets ($P < 0.05$). Meishan piglets also had lower concentrations of TG and LDL-C, although this difference did not reach statistical significance. Other parameters were not significantly affected by breed.

3.2. Comparisons of digestive enzyme activities between Meishan and Yorkshire piglets

Meishan piglets had higher activity of gastric lipase but lower activity of jejunal lactase than Yorkshire piglets ($P < 0.05$) (Table 2). Activity of jejunal amylase in Meishan piglets trended to be lower than that in Yorkshire piglets ($P = 0.09$). There was no valid data of pepsin and jejunal protease activities, which were under detection level of the enzyme kits.

3.3. Comparisons of VFA concentrations between Meishan and Yorkshire piglets

Concentrations of acetate, propionate and total VFA in the cecum contents of Meishan piglets were significantly higher than those of Yorkshire piglets, while acetate concentration in jejunum content was lower in Meishan piglets than in Yorkshire piglets ($P < 0.05$) (Table 3). Molar percentages of acetate, propionate and butyrate in the cecum, colon and feces samples of the two breeds are presented in Table 4. Compared with Yorkshire piglets, Meishan piglets had a higher proportion of propionate in the cecum and colon, but a lower acetate:propionate ratio in the colon ($P < 0.05$). In jejunum, no VFA were detected except for acetate because of their extremely low concentrations.

Table 1

Blood biochemical parameters of Meishan and Yorkshire piglets.

Item	Meishan piglets	Yorkshire piglets	<i>P</i> -value
TP, g/L	58.20 \pm 6.19	61.38 \pm 4.61	0.689
ALB, g/L	37.38 \pm 3.71	36.80 \pm 1.86	0.891
GLO, g/L	20.82 \pm 3.18	23.75 \pm 3.01	0.518
GLU, mmol/L	7.70 \pm 0.74	8.93 \pm 0.58	0.218
Urea, mmol/L	4.52 \pm 0.87	3.65 \pm 0.19	0.351
TG, mmol/L	0.91 \pm 0.18	1.49 \pm 0.30	0.129
HDL-C, mmol/L	1.53 \pm 0.09 ^b	2.28 \pm 0.27 ^a	0.036
LDL-C, mmol/L	2.14 \pm 0.35	3.15 \pm 0.58	0.166
TC, mmol/L	3.57 \pm 0.33 ^b	5.45 \pm 0.62 ^a	0.023

TP = total protein; ALB = albumin; GLO = globulin; GLU = glucose; TG = triglycerides; HDL-C = high-density lipoprotein cholesterol; LDL-C = low-density lipoprotein cholesterol; TC = total cholesterol.

^{a,b} Within a row, means without a common letter differ ($P < 0.05$).

Download English Version:

<https://daneshyari.com/en/article/4522296>

Download Persian Version:

<https://daneshyari.com/article/4522296>

[Daneshyari.com](https://daneshyari.com)