



Short communication

β -Cyclodextrin-based nanosponges improve 1-MCP efficacy in extending the postharvest quality of cut flowers

Ludovica Seglie^a, Marco Devecchi^a, Francesco Trotta^b, Valentina Scariot^{a,*}^a Department of Agricultural, Forest and Food Sciences, University of Torino, Via Leonardo da Vinci 44, 10095 Grugliasco, TO, Italy^b Department of Chemistry, University of Torino, Via Pietro Giuria 7, 10125 Turin, Italy

ARTICLE INFO

Article history:

Received 9 January 2013

Received in revised form 8 May 2013

Accepted 16 May 2013

Keywords:

Ethylene inhibitor

Flower longevity

Nano-carrier

Preservative

Senescence

Vase life

ABSTRACT

Investigations of the effect of the β -cyclodextrin-based nanosponge-1-methylcyclopropene (1-MCP) complex (β -CD-NS complex) on six ethylene-sensitive cut flowers were performed. *Anemone coronaria* L. multicolor, *Ranunculus asiaticus* L. 'Minou Abrown', *Helianthus annuus* L. 'Sunrich Orange', *Rosa hybrida* L. 'Jupiter', *Paeonia lactiflora* L. 'Sarah Bernhardt', and *Papaver nudicaule* L. multicolor underwent four different treatments: a vase placement in a suspension of β -CD-NS complex ($0.25 \mu\text{L L}^{-1}$), exposure to gaseous 1-MCP ($0.25 \mu\text{L L}^{-1}$) for 6 h, exposure to exogenous ethylene ($1 \pm 0.2 \mu\text{L L}^{-1}$) for 24 h, and placement in tap water.

β -Cyclodextrin-based nanosponges proved to enhance 1-MCP efficacy in all the tested species. According to the species, senescence symptoms were decreased or delayed. Anemone and poppy had a longer esthetic flower quality and vase life, and a reduced petal abscission. In ranunculus, β -CD-NS complex improved longevity. In sunflower, it also maintained longer chlorophyll leaf content and cut flower ornamental value. Peony opened more slowly and rose maintained its ornamental quality longer than 1-MCP treated flowers. This study opens possibilities for commercial use of the β -CD-NS complex in the floriculture industry.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Postharvest performance is of crucial importance to the value of cut flowers. The life span of flowers is generally determined by the time to abscission of petals that are still turgid, or by the time to petal wilting or withering. In many species, these phenomena are regulated by ethylene (Woltering and van Doorn, 1988; van Doorn, 2001). Premature senescence and abscission resulting from exposure to exogenous or endogenous ethylene can be controlled several ways including ethylene action inhibitors (Martínez-Romero et al., 2007) quite commonly.

In particular, while 1-methylcyclopropene (1-MCP), has proved itself to be a very effective preventer of negative ethylene responses, its gaseous nature makes treatment difficult (Serek and Sisler, 2005; Serek et al., 2006).

Recently, we developed a non-volatile formulation of 1-MCP in a β -cyclodextrin-based nanosponge (β -CD-NS 1:8) structure. Supplied in the conservation solution, this formulation (β -CD-NS

complex) prolonged the vase life of carnation cut flowers (Seglie et al., 2011a,b), and controlled *Botrytis cinerea* damage (Seglie et al., 2012) better than gaseous 1-MCP.

While carnation is generally used in studies on postharvest preservation of ethylene-sensitive flowers (Woltering and van Doorn, 1988; Serek et al., 1995a,b); it is not a universal paradigm for flower senescence. The response to ethylene varies widely according to species (Reid and Wu, 1992). Therefore, our study evaluated the effectiveness of the β -CD-NS complex to improve cut flower vase life on a number of ethylene-sensitive species.

2. Materials and methods

The flower market of Sanremo (UCFlor, Liguria, Italy) seasonally provided cut flowers of *Anemone coronaria* L. multicolor, *Ranunculus asiaticus* L. 'Minou Abrown', *Helianthus annuus* L. 'Sunrich Orange', *Rosa hybrida* L. 'Jupiter', *Paeonia lactiflora* Pall. 'Sarah Bernhardt', and *Papaver nudicaule* L. multicolor.

Stems were transferred to the postharvest laboratory of the Department of Agricultural, Forest and Food Sciences at the University of Turin. Experimental evaluations of vase life were performed in an imitated inside environment (IE) maintained at $20 \pm 2^\circ\text{C}$, 60% RH, and 12 h daily of $46 \mu\text{mol m}^{-2} \text{s}^{-1}$ cool white light (meter model HT307; HT, Faenza, Italy).

Abbreviations: CD, cyclodextrin; 1-MCP, 1-methylcyclopropene; β -CD-NS, β -CD-based-nanosponge; β -CD-NS complex, 1-MCP included in β -CD-NS.

* Corresponding author. Tel.: +39 011 6708932; fax: +39 011 6708798.

E-mail address: valentina.scariot@unito.it (V. Scariot).

Cut flowers were re-cut, labeled and re-watered. Each species sample was divided into sub-groupings of six cut stems (each 30 cm long after re-cutting). Each sub-group was then treated according to the following schemes: in vases with an aqueous suspension of β -CD-NS complex to supply $0.25 \mu\text{L L}^{-1}$ of active ingredient; in an air tight cabinet (112L) exposed to equal concentrations of volatile 1-MCP (3.3% a. i., SmartFresh™, AgroFresh Inc., USA) for 6 h; in another air tight cabinet exposed to exogenous ethylene ($1 \pm 0.2 \mu\text{L L}^{-1}$) for 24 h; and in vases of tap water as a control. The experiment was performed twice for each species.

Endogenous ethylene production of the control flowers was measured using a digital Agilent Technologies gas chromatograph, 7890A Network GC system (Santa Clara, California). N_2 at 40 mL min^{-1} was used as the gas carrier at a column temperature 60°C and calibration range between 0.5 and 3 ppm. For each species, six samples were considered.

Flower postharvest performance was measured and visual checks for symptoms of variation in senescence were performed daily. In particular, we monitored senescence level (1=initial senescence, 2=complete senescence; Seglie et al., 2011a), flower opening stage (1=initial opening, 2=half opening, 3=full opening; Guo et al., 2004, modified), and abscission. Chlorophyll content was indirectly measured in leaves through the Chlorophyll Meter SPAD-502 instrument (Konica Minolta Sensing Inc., Osaka, Japan).

Analysis of variance (ANOVA) was performed to assess statistical significance among mean values using the Ryan–Einot–Gabriel–Welsch's multiple step-down F (REGW-F) test ($P \leq 0.05$) with SPSS Software (Chicago, USA).

3. Results and discussion

Senescence performance and ethylene sensitivity are highly species related (van Doorn, 2001). Large variabilities have been observed even among cultivars, such as the Asian buttercup (Kenza et al., 2000; Scariot et al., 2009), rose (Chanami et al., 2005), and peony (Hoffman et al., 2010; Eason et al., 2002). Flowers that respond to low concentrations of exogenous ethylene are probably those in which ethylene is naturally involved in senescence (Reid and Wu, 1992). In this study, ethylene hastened the symptoms of natural senescence in all species. However, a climacteric pattern was observed only in poppy (ethylene up to $12 \mu\text{L L}^{-1}$ at day 5) while in peony and sunflower ethylene production was not detected (data not shown).

Anemone cut flowers lose their decorative value in a very short period. Pedicels elongate excessively and petals wilt, lose their color, become translucent, and abscise (Meir et al., 2007). Previous studies have already demonstrated the ethylene sensitivity of the anemone (Armitage and Laushman, 1993) and the efficacy of ethylene antagonists (silver nitrate and silver thiosulphate, STS) in extending its vase life and preventing petal abscission (Sharifani et al., 2005). In this study, the application of β -CD-NS complex was found to prolong cut flower esthetic quality and longevity for up to 1.5 additional days (Fig. 1), slow stages 1 and 2 of flower opening (Table 1), and completely prevent abscission (Fig. 2) more effectively than gaseous 1-MCP. Petal rolling over and stem folding (Sharifani et al., 2005), which usually occur in anemone cut flowers, were also limited.

Asian buttercup flower senescence is characterized by loss of turgor and petal color change followed by petal wilting and drop when lightly touched. Previous tests of the effects of silver thiosulphate (STS) and aminooxyacetic acid (AOA) on buttercup postharvest performance have yielded unsatisfactory results in enhancing vase life (Shari et al., 2011). In this study, ethylene antagonists were effective in delaying senescence symptoms with β -CD-NS complex; they outperformed gaseous 1-MCP as




Fig. 1. Longevity of six species of cut flowers according to four treatments: placement in a vase suspension of β -CD-NS complex ($0.25 \mu\text{L L}^{-1}$), exposure to gaseous 1-MCP ($0.25 \mu\text{L L}^{-1}$) for 6 h, exposure to exogenous ethylene ($1 \pm 0.2 \mu\text{L L}^{-1}$) for 24 h, and placement in tap water. The count of days to reach the initial stage of senescence (Level 1, line) and complete senescence (Level 2, dotted line) was measured. Mean separation within bars by the Ryan–Einot–Gabriel–Welsch's multiple stepdown F (REGW-F) test, $P \leq 0.001$.

demonstrated by delay of complete senescence (9.7 versus 8 days; Fig. 1) and original flower color maintenance (data not shown).

Previous studies of sunflower have shown exogenous ethylene applications result in short vase life, flower abscission (Redman et al., 2002), and in largely, wilted basal foliage immediately after anthesis (Pallez et al., 2002; Castro et al., 2011). Furthermore, ethylene antagonist treatment has failed to limit or delay senescence (Redman et al., 2002). In this study, chlorophyll degradation

Table 1

Flower opening of six flower species according to four treatments: placement in a vase suspension of β -CD-NS complex ($0.25 \mu\text{L L}^{-1}$), exposure to a gaseous 1-MCP ($0.25 \mu\text{L L}^{-1}$) for 6 h, exposure to exogenous ethylene ($1 \pm 0.2 \mu\text{L L}^{-1}$) for 24 h, and placement in tap water. Stages from 1 to 3 (1=initial opening, 2=half opening, 3=full opening), and the duration (in day) of each stage were considered.

Species	Treatments	Flower opening (days)		
		Stage 1	Stage 2	Stage 3
<i>A. coronaria</i>	β -CD-NS complex	2.17a [*]	2.00a	2.50a
	1-MCP	1.50b	1.50b	3.67a
	Ethylene	0.17c	0.50d	1.50b
	Water	0.33c	1.00c	1.33b
<i>R. asiaticus</i>	β -CD-NS complex	0.17	2.83a	2.67a
	1-MCP	0.67	2.50a	2.83a
	Ethylene	0.83	1.17b	1.00b
	Water	0.83	1.33b	2.00b
<i>H. annuus</i>	β -CD-NS complex	4.00a	3.17a	2.83a
	1-MCP	2.83b	2.67b	2.17a
	Ethylene	0.33d	2.00c	1.00bc
	Water	1.00c	2.67b	1.50b
<i>R. hybrida</i>	β -CD-NS complex	1.00	3.67a	3.33a
	1-MCP	1.00	2.50b	2.67b
	Ethylene	0.33	1.83c	1.33d
	Water	0.33	2.17bc	2.17c
<i>P. lactiflora</i>	β -CD-NS complex	2.83a	3.33a	2.67a
	1-MCP	1.50b	2.67b	2.00b
	Ethylene	0.50c	1.00c	0.83d
	Water	0.67c	1.83c	1.33c
<i>P. nudicaule</i>	β -CD-NS complex	2.67a	2.33a	2.50a
	1-MCP	2.00b	1.83b	1.33b
	Ethylene	0.50c	0.83c	1.00c
	Water	0.67c	1.33bc	1.17bc

^{*} Mean separation within columns and stages by the Ryan–Einot–Gabriel–Welsch's multiple stepdown F (REGW-F) test, $P \leq 0.001$.

Download English Version:

<https://daneshyari.com/en/article/4567235>

Download Persian Version:

<https://daneshyari.com/article/4567235>

[Daneshyari.com](https://daneshyari.com)