

Assessment of morphological changes and determination of best cane collection time for 140 RU and 5 BB

Alper Dardeniz^{a,*}, Nuray Mücellâ Müftüoğlu^b, Zeliha Gökbayrak^a, Mustafa Fırat¹

^aÇanakkale Onsekiz Mart University, Faculty of Agriculture, Department of Horticulture, 17020 Çanakkale, Turkey

^bÇanakkale Onsekiz Mart University, Faculty of Agriculture, Department of Soil, 17020 Çanakkale, Turkey

Received 24 February 2006; received in revised form 22 January 2007; accepted 30 January 2007

Abstract

This study was carried out to assess morphological changes in the canes of 140 Ru and 5 BB and to determine best time for cane collection. The first part of the study included the morphological assessment of the rootstock canes during the winters of 2003–2004 and 2004–2005. Canes collected in the second part of the study in 2005–2006 were grown in pots to relate the changes with the vegetative growth characteristics.

Cane collection commenced at leaf fall and they were collected three more times at 15-day intervals. Morphological changes were determined from the transverse internodal cuts and included: width of pith, xylem, phloem plus bark tissues, ratios of cane width/pith, cane width/bark + phloem, xylem/pith and xylem/bark + phloem, and cane water content. Findings showed that lignification was best 30 and 45 days after leaf fall for 140 Ru and 15, 30 and 45 days after leaf fall for 5 BB.

Canes collected in 2005–2006 were used to relate the changes observed in the previous 2 years with the vegetative development. Two-bud cuttings were grown in pots and viable plant and rooting ratios, root fresh weight, shoot number and length, and node number were determined. Data obtained were found in agreement with the morphological findings. A 140 Ru had the best vegetative growth 30 and 45 days after leaf fall, and 5 BB grew well at each collection period.

Results showed that it is best not to collect canes from rootstock nurseries at or right after leaf fall due to poor lignification and subsequent poor growth.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Grape rootstock; Cane; Morphology; Lignification; Vegetative growth

1. Introduction

Phylloxera (*Daktulosphaira vitifoliae* Fitch.) is one of the most significant limiting factors to grape-growing. The only proven method for long term control is to graft susceptible *Vitis vinifera* onto phylloxera tolerant vines, rootstocks, developed from native American grapevine species (Granett et al., 2001).

Cuttings well lignified and rich with reserves are better at rooting and growth than poor ones (Gautheret, 1966; Thorpe, 1974; Kismali, 1981; Bartolini et al., 1996; Hunter et al., 2004). In addition, callusing is higher when cuttings are highly lignified (Kozma et al., 1972; Kismali, 1978). Due to the high work load in large nurseries during winter, rootstock cuttings are usually started collecting right after leaf fall (end of

November) and stored in cold rooms until grafting. Collecting rootstock canes at leaf fall is however not advised since translocation, conversion and storage of reserves continue after defoliation in the over wintering parts of vines (Coombe and Dry, 1992; Çelik, 1998) even if only at a very low level.

Wood fibers, wood parenchyma and ray regions are important sugar and starch storage organs (Eifert et al., 1961). Conversion from sugar to starch or vice versa takes place depending on winter air temperature (Winkler et al., 1974). Starch is at maximum and sugars are at minimum level in December and January. Conversion of sugars to starch starts in spring, but the overall reserve carbohydrate level declines due to consumption by the breaking buds (Eifert et al., 1961; Coombe and Dry, 1992). Harvesting of dormant cuttings is therefore advised for mid winter and not at the beginning of winter or when vines start taking up water in the spring (Kocamaz, 1995; Çelik, 1998).

The degree of lignification is best assessed by using the ratios of cane width/pith and phloem + xylem/pith values

* Corresponding author.

E-mail address: adardeniz@comu.edu.tr (A. Dardeniz).

¹ Agricultural Engineer.

(Oraman, 1963; Dardeniz, 2001). There are different factors affecting the cane width/pith ratio. Dardeniz (2001) studied the effect of cluster thinning on cane lignification. He found that 30–60% cluster thinning increased cane width/pith ratio in Cardinal and Amasya compared to the controls. Dardeniz and Kismalı (2001) found that the cane width/pith ratio of 140 Ru decreased with increasing shoot number. Dardeniz and Şahin (2005) examined cane hardening on the canes of grape cultivars Yalova İncisi (*Vitis vinifera* L.) grafted onto different rootstocks. Yalova İncisi had the best hardening on 1103 P, followed by 5 BB, 41 B and 140 Ru. Lignification level (cane width/pith) was determined on rootstock 140 Ru and 1103 P with different shoot loads (4, 8, 12 shoots per vine) (Dardeniz and Kismalı, 2001). 140 Ru had the highest levels at four and eight shoot/vine. Best cane width/pith ratio was obtained from four shoots per vine for 1103 P. The same shoots were used as 2-bud cuttings in the winter and were grown in pots. It was found that shoot weight, shoot length and callus weight increased significantly as the cane matured.

Pith in 1-year-old canes has short-thin walled cells with amorphous shapes and it contains chlorophyll pigments (Oraman, 1972; Çelik, 1998; Ağaoğlu, 1999). Properly grown and matured canes are well lignified when their piths are narrow and xylem tissues are wide (Çelik, 1998). Pith is much wider in young shoots, decreasing in width as they develop during the season (Ağaoğlu, 1999).

The morphological changes that occur in shoots of American grape species rootstocks at or right after leaf fall have not been fully studied. These kinds of detailed studies will enable both researchers and nurserymen to determine when the pith is narrow or the ratios of cane width/pith and xylem/pith are high to select the appropriate time for harvesting canes. The objective of this study was to periodically determine morphological changes in the canes of the rootstocks and correlate this data to vegetative development of the rootstocks.

2. Material and method

This study consisted of two parts. The first part comprised observations on the morphological changes in the canes of grapevine rootstocks, 140 Ru (*Vitis berlandieri* x *V. rupestris*) and 5 BB (*V. berlandieri* x *V. riparia*). Canes were collected in the winters of 2003–2004 and 2004–2005 from the Fruit Propagation Station, located in Umurbey, Çanakkale, Turkey. In the second part (winter season of 2005–2006), rootstock canes were grown in pots to collect data on vegetative growth characteristics and relate these with the findings obtained in the first part.




Fig. 1. Cut places on the canes of the rootstocks.

Rootstocks in the foundation block were 14-year-old, growing on the ground with 2 m × 2 m spacing. They were grown under dry land conditions with normal cultivation and fertilization. No shoot management techniques were applied during the study. Table 1 presents the soil characteristics of the block.

The trial was designed in randomized plots comprising four replicates and three vines per replicate. Canes were collected four times, leaf fall (20–25 November, 80% defoliation), 15 days later (5–10 December), 30 days later (20–25 December), and 45 days later (5–10 January). Canes were randomly chosen on the basis of equal thickness of the canes at each and every collection time. Two canes per vine were selected and cut into 4-bud pieces (1–4, 5–8, 9–12, 13–16, 17–20) to facilitate morphological measurements. The pieces were cut in the middle of the internodes (Fig. 1). The number of the transverse cuts per rootstock per year was 1440 (3 vines × 2 canes × 15 cuts × 4 replicates × 4 times). Characteristics listed below were measured with the aid of digital compass at the cut surfaces as shown in Fig. 2.

Cane width (mm): the arithmetic mean of the thinnest (CW_1) and the thickest (CW_2) points of the cane, xylem width (mm); the arithmetic mean of four corresponding point measurements at cut surface (X_1, X_2, X_3, X_4), pith width (mm); the arithmetic mean of the thinnest (P_1) and thickest (P_2) points of the cane, bark + phloem width (mm); the arithmetic mean of the visually determined thinnest (BP_1) and thickest (BP_2) points of the cane, water content (%) as described by (Kacar, 1972); [(fresh weight – dry weight) × 100/fresh weight]. The ratios of cane width/pith and xylem/pith were calculated to determine the lignification levels. Cane width/bark + phloem and xylem/bark + phloem ratios were also calculated for wood hardening determination.

The objective of the second part of study was to compare and relate the first part findings with the vegetative growth of the canes. The canes were collected from the same vines in the foundation block. Canes were also cut into 4-bud segments (1–4, 5–8, 9–12, 13–16, 17–20) to ease handling. They were treated with a fungicide (Captan, Southern Agricultural

Table 1
Soil profile features of the rootstock foundation block

Depth (cm)	Texture	Salinity (%)	pH (1:2.5)	Calcareous (CaCO ₃ %)	P (kg day ⁻¹ P ₂ O ₅)	K (kg day ⁻¹ K ₂ O)	Organic matter (%)
0–30	Loamy	None	7.5–8.0	Low	High	Adequate	Low
30–60	Loamy	None	7.5–8.0	High	Low	Adequate	Low
60–90	Loamy	None	7.5–8.0	High	Low	Adequate	Very low

Download English Version:

<https://daneshyari.com/en/article/4569707>

Download Persian Version:

<https://daneshyari.com/article/4569707>

[Daneshyari.com](https://daneshyari.com)