



Radial eigenvectors of the Laplacian of the nonbinary hypercube

Murali K. Srinivasan

Department of Mathematics, Indian Institute of Technology, Bombay, Powai, Mumbai 400 076, India

ARTICLE INFO

Article history:

Received 15 August 2012

Accepted 6 October 2012

Available online 29 November 2012

Submitted by R.A. Brualdi

AMS classification:

05C50

Keywords:

Laplacian eigenvalues

Nonbinary hypercube

Radial eigenvectors

ABSTRACT

Let q be a positive integer ≥ 2 . Define a $(n+1) \times (n+1)$ real, symmetric, tridiagonal matrix M , with rows and columns indexed by $\{0, 1, \dots, n\}$, and with entries given by:

$$M(i, j) = \begin{cases} -\sqrt{(q-1)j(n-j+1)} & \text{if } i = j-1, \\ j + (q-1)(n-j) & \text{if } i = j, \\ -\sqrt{(q-1)(j+1)(n-j)} & \text{if } i = j+1, \\ 0 & \text{if } |i-j| \geq 2. \end{cases}$$

The $(n+1)$ -dimensional space of radial vectors for the nonbinary hypercube $C_q(n)$ is invariant under the Laplacian and M arises as the matrix of the Laplacian with respect to a suitable orthonormal basis of radial vectors. We show that the eigenvalues of M are $0, q, 2q, \dots, nq$ by explicitly writing down the eigenvectors.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

For a finite set S we denote by $V(S)$ the complex vector space with S as basis. For a simple graph G on the vertex set S , recall that the *adjacency operator* $A : V(S) \rightarrow V(S)$ of G is defined, for $v \in S$, by $A(v) = \sum_u u$, where the sum is over all vertices u adjacent to v .

Let $B(n) = \{a = (a_1, \dots, a_n) : a_i = 0 \text{ or } 1 \text{ for all } i\}$ denote the set of all sequences of 0's and 1's of length n . The *binary hypercube* $C(n)$ is the simple graph with $B(n)$ as the vertex set and with $a, b \in B(n)$ connected by an edge provided they differ in exactly one coordinate.

E-mail addresses: mks@math.iitb.ac.in, murali.k.srinivasan@gmail.com

The spectrum of the adjacency operator A of $C(n)$ is easily determined using the graph operation of Cartesian products (see Section 1.4.6 in [1] or Section 2.5 in [3] where this operation is called sum). We recall this briefly (without proof). First we find that the adjacency eigenvalues of $C(1)$ are $\alpha = 1$ and $\beta = -1$. Now $C(n)$ is the n -fold Cartesian product of $C(1)$ and thus the eigenvalues of A are $(n-i)\alpha + i\beta = n-2i$ with multiplicity $\binom{n}{i}$, for $i = 0, 1, \dots, n$. This method also gives an explicit set of eigenvectors (using tensor products of adjacency eigenvectors of $C(1)$) of A .

In the theory of lumped Markov chains the following subspace of $V(B(n))$, invariant under A , is considered. Define the following vectors in $V(B(n))$:

$$u_i = \sum_a a, \quad i = 0, 1, \dots, n, \quad (1)$$

where the sum is over all $a = (a_1, \dots, a_n) \in B(n)$ containing i 1's. Let $R(n)$ be the subspace of $V(B(n))$ spanned by $\{u_0, \dots, u_n\}$. Clearly, $\dim(R(n)) = n+1$. Elements of $R(n)$ are called *radial vectors* in $V(B(n))$. It is easy to see that $R(n)$ is invariant under A . We have

Theorem 1.1. *The eigenvalues of $A : R(n) \rightarrow R(n)$ are $n-2i$, $i = 0, 1, \dots, n$ and the (canonically defined) eigenvectors can be explicitly written down.*

Theorem 1.1 is proved, in an equivalent form, in [2] (see Theorem 2.6.4) in the context of radial harmonic analysis on the binary hypercube. In this note we present a nonbinary generalization of Theorem 1.1. The main point of this generalization is that the correct formulation of the nonbinary analog, so as to obtain integral eigenvalues, involves the Laplacian operator of a nonregular graph. In the binary case the adjacency and Laplacian formulations are equivalent due to the regularity of $C(n)$.

2. Nonbinary hypercube

Let $q \geq 2$ be a positive integer and define

$$B_q(n) = \{(a_1, \dots, a_n) : a_i \in \{0, 1, \dots, q-1\} \text{ for all } i\}.$$

For $a \in B_q(n)$ define the *support* of a by $S(a) = \{i \in \{1, 2, \dots, n\} : a_i \neq 0\}$. We define the *nonbinary hypercube* $C_q(n)$ to be the simple graph with $B_q(n)$ as the vertex set and with $a, b \in B_q(n)$ connected by an edge provided $S(a) \subseteq S(b)$ or $S(b) \subseteq S(a)$, $||S(a)| - |S(b)|| = 1$, and $a_i = b_i$, for $i \in S(a) \cap S(b)$. That is, a and b are connected by an edge provided they differ in exactly one coordinate and one of a or b is 0 in that coordinate. The degree of a vertex a of $C_q(n)$ is $|S(a)| + (q-1)(n - |S(a)|)$. Note that, while $C(n) = C_2(n)$ is a regular graph (of degree n), $C_q(n)$ is not regular for $q \geq 3$. Note also that $C_q(n)$ is different from what is usually called the nonbinary Hamming graph on q letters (which is regular).

Let A_q denote the adjacency operator of $C_q(n)$ and let $\deg : V(B_q(n)) \rightarrow V(B_q(n))$ denote the *degree operator* given by: $\deg(a) = (\text{degree of } a) a$, $a \in B_q(n)$. Recall that the *Laplacian operator* L_q of $C_q(n)$ is defined by $L_q = \deg - A_q$.

Define the following vectors in $V(B_q(n))$:

$$v_i = \sum_a a, \quad i = 0, 1, \dots, n, \quad (2)$$

where the sum is over all $a \in B_q(n)$ with $|S(a)| = i$. Let $R_q(n)$ be the subspace of $V(B_q(n))$ spanned by $\{v_0, \dots, v_n\}$. Clearly, $\dim(R_q(n)) = n+1$. Elements of $R_q(n)$ are called *radial vectors* in $V(B_q(n))$. It is easy to see that $R_q(n)$ is invariant under L_q . Before determining the eigenvalues of $L_q : R_q(n) \rightarrow R_q(n)$ we briefly discuss the entire Laplacian spectrum of $C_q(n)$, i.e., the eigenvalues of $L_q : V(B_q(n)) \rightarrow V(B_q(n))$. First we find that the Laplacian spectrum of $C_q(1)$ is $\{0, q, 1\}$, with 0 and q of multiplicity 1, and 1 of multiplicity $q-2$. Now $C_q(n)$ is the n -fold Cartesian product of $C_q(1)$ and thus its Laplacian

Download English Version:

<https://daneshyari.com/en/article/4600590>

Download Persian Version:

<https://daneshyari.com/article/4600590>

[Daneshyari.com](https://daneshyari.com)