



# A cotangent sum related to zeros of the Estermann zeta function



Michael Th. Rassias

Department of Mathematics, ETH-Zürich, Rämistrasse 101, 8092 Zürich, Switzerland

## ARTICLE INFO

### Keywords:

Cotangent sums  
Asymptotic approximation  
Taylor expansion  
Estermann zeta function  
Riemann hypothesis  
Fractional part

## ABSTRACT

We consider a cotangent sum related to Estermann's zeta function. We provide an elementary and self-contained improvement of the error term in an asymptotic formula proved by V.I. Vasyunin.

© 2014 Elsevier Inc. All rights reserved.

## 1. Introduction

The problem of the computation of zeros of the Estermann zeta function at  $s = 0$  reduces to the calculation of a specific cotangent sum. However, before we proceed with the presentation of our result related to that cotangent sum, we will exhibit a general framework which motivated the research in this area and explain how this problem is related to fundamental open problems, such as the Riemann hypothesis.

The Estermann zeta function  $E(s, \frac{h}{k}, \alpha)$  is defined by the Dirichlet series

$$E\left(s, \frac{h}{k}, \alpha\right) = \sum_{n \geq 1} \frac{\sigma_{\alpha}(n) \exp(2\pi i h n / k)}{n^s},$$

where  $h, k$  are positive integers,  $(h, k) = 1$ ,  $\text{Re } s > \max(1, 1 + \text{Re } \alpha)$ ,  $k \geq 1$  and

$$\sigma_{\alpha}(n) = \sum_{d|n} d^{\alpha}.$$

It is a known fact (see [4,7]) that the Estermann zeta function can be continued analytically to a meromorphic function, on the whole complex plane up to two simple poles  $s = 1$  and  $s = 1 + \alpha$  if  $\alpha \neq 0$  or a double pole at  $s = 1$  if  $\alpha = 0$ .

In 1995, Ishibashi (see [6]) presented a nice result concerning the value of  $E(s, \frac{h}{k}, \alpha)$  at  $s = 0$ .

**Theorem 1.1** (Ishibashi). *Let  $k \geq 2$ ,  $1 \leq h < k$ ,  $(h, k) = 1$ ,  $\alpha \in \mathbb{N} \cup \{0\}$ . Then*

(1) *For even  $\alpha$ , it holds*

$$E\left(0, \frac{h}{k}, \alpha\right) = \left(-\frac{i}{2}\right)^{\alpha+1} \sum_{m=1}^{k-1} \frac{m}{k} \cot^{(\alpha)}\left(\frac{\pi m h}{k}\right) + \frac{1}{4} \delta_{\alpha,0},$$

where  $\delta_{\alpha,0}$  is the Kronecker delta function.

E-mail address: [michail.rassias@math.ethz.ch](mailto:michail.rassias@math.ethz.ch)

<http://dx.doi.org/10.1016/j.amc.2014.04.086>

0096-3003/© 2014 Elsevier Inc. All rights reserved.

(2) For odd  $\alpha$ , it holds

$$E\left(0, \frac{h}{k}, \alpha\right) = \frac{B_{\alpha+1}}{2(\alpha+1)}.$$

In the case when  $k = 1$  we have

$$E(0, 1, \alpha) = \frac{(-1)^{\alpha+1} B_{\alpha+1}}{2(\alpha+1)},$$

where by  $B_m$  we denote the  $m$ th Bernoulli number.

Hence for  $k \geq 2, 1 \leq h < k, (h, k) = 1$ , it follows that

$$E\left(0, \frac{h}{k}, 0\right) = \frac{1}{4} + \frac{i}{2} c_0\left(\frac{h}{k}\right),$$

where

$$c_0\left(\frac{h}{k}\right) = -\sum_{m=1}^{k-1} \frac{m}{k} \cot\left(\frac{\pi m h}{k}\right).$$

This is exactly the cotangent sum that we are going to investigate in this paper. A very natural question that rises, is why this specific form of the Estermann zeta function is interesting. One example which demonstrates its importance is the following.

In 1985, Balasubramanian et al. (see [2]), used properties of  $E(s, \frac{h}{k}, \alpha)$  to prove an asymptotic formula for

$$I = \int_0^T \left| \zeta\left(\frac{1}{2} + it\right) \right|^2 \left| A\left(\frac{1}{2} + it\right) \right|^2 dt,$$

where  $A(s)$  is a Dirichlet polynomial.

Asymptotics for functions of the form of  $I$  are useful for theorems which provide a lower bound for the portion of zeros of the Riemann zeta-function on the critical line (see [8,9]). Period functions and families of cotangent sums appear in recent work of Bettin and Conrey (see [3]). They generalize the Dedekind sum and share with it the property of satisfying a reciprocity formula. They have proved a reciprocity formula for the Vasyunin sum, which appears in the Nyman–Beurling criterion (see [1]) for the Riemann hypothesis.

In 1995 Vasyunin (see [10]), using Riemann sums, proved in a short elegant way the following theorem.

**Theorem 1.2** (Vasyunin). For large integer values of  $b$ , we have

$$c_0\left(\frac{1}{b}\right) = \frac{1}{\pi} b \log b - \frac{b}{\pi} (\log 2\pi - \gamma) + O(\log b),$$

where  $\gamma$  is the Euler–Mascheroni constant.

In this paper we provide an elementary and self-contained improvement of the error term in Vasyunin’s asymptotic formula. Namely, we prove the following.

**Theorem 1.3.** For large integer values of  $b$ , we have

$$c_0\left(\frac{1}{b}\right) = \frac{1}{\pi} b \log b - \frac{b}{\pi} (\log 2\pi - \gamma) + O(1).$$

## 2. Construction and some approximations for $c_0(1/b)$

It is evident that for  $a, b \in \mathbb{N}, b \geq 2$ , the integer part  $[a/b]$  is equal to the number of integers between 1 and  $a$  which are divisible by  $b$ . However, it is a basic fact that

$$\frac{1}{b} \sum_{m=0}^{b-1} e^{2\pi i m k/b} = \begin{cases} 1, & \text{if } b \mid k \\ 0, & \text{otherwise.} \end{cases}$$

Hence, it follows that

$$\left[ \frac{a}{b} \right] = \frac{1}{b} \sum_{k=1}^a \sum_{m=0}^{b-1} e^{2\pi i m k/b}.$$

Download English Version:

<https://daneshyari.com/en/article/4627397>

Download Persian Version:

<https://daneshyari.com/article/4627397>

[Daneshyari.com](https://daneshyari.com)