



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Combinatorics

Bounds on the vertex–edge domination number of a tree



Bornes sur le nombre de domination sommet–arête d'un arbre

Balakrishna Krishnakumari^a, Yanamandram B. Venkatakrishnan^a,
Marcin Krzywkowski^{b,c,1,2}

^a Department of Mathematics, SASTRA University, Thanjavur, Tamil Nadu, India

^b Department of Mathematics, University of Johannesburg, South Africa

^c Faculty of Electronics, Telecommunications and Informatics, Gdansk University of Technology, Poland

ARTICLE INFO

Article history:

Received 22 November 2013

Accepted after revision 18 March 2014

Available online 3 April 2014

Presented by the Editorial Board

ABSTRACT

A vertex–edge dominating set of a graph G is a set D of vertices of G such that every edge of G is incident with a vertex of D or a vertex adjacent to a vertex of D . The vertex–edge domination number of a graph G , denoted by $\gamma_{ve}(T)$, is the minimum cardinality of a vertex–edge dominating set of G . We prove that for every tree T of order $n \geq 3$ with l leaves and s support vertices, we have $(n - l - s + 3)/4 \leq \gamma_{ve}(T) \leq n/3$, and we characterize the trees attaining each of the bounds.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

RÉSUMÉ

Un ensemble sommet–arête dominant d'un graphe G est un ensemble D de sommets de G tel que chaque arête de G soit incidente à un sommet de D ou à un sommet adjacent à un sommet de D . Le nombre de domination sommet–arête d'un graphe G , noté $\gamma_{ve}(T)$, est le cardinal minimum d'un ensemble sommet–arête dominant de G . Nous prouvons que, pour chaque arbre T d'ordre $n \geq 3$ avec l feuilles et des sommets s de soutien, que nous avons $(n - l - s + 3)/4 \leq \gamma_{ve}(T) \leq n/3$, et nous caractérisons les arbres atteignant chacune des limites.

© 2014 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

Let $G = (V, E)$ be a graph. By the neighborhood of a vertex v of G we mean the set $N_G(v) = \{u \in V(G) : uv \in E(G)\}$. The degree of a vertex v , denoted by $d_G(v)$, is the cardinality of its neighborhood. By a leaf we mean a vertex of degree one, while a support vertex is a vertex adjacent to a leaf. We say that a support vertex is strong (weak, respectively) if it is adjacent to at least two leaves (exactly one leaf, respectively). We denote by P_n the path on n vertices. Let T be a tree, and let v be a vertex of T . We say that v is adjacent to a path P_n if there is a neighbor of v , say x , such that the subtree

E-mail address: marcin.krzywkowski@gmail.com (M. Krzywkowski).

¹ Research fellow at the Department of Mathematics, University of Johannesburg, South Africa.

² Research partially supported by the Polish National Science Centre grant 2011/02/A/ST6/00201.

resulting from T by removing the edge vx and which contains the vertex x as a leaf, is a path P_n . By a star we mean a connected graph in which exactly one vertex has degree greater than one.

A subset $D \subseteq V(G)$ is a dominating set of G if every vertex of $V(G) \setminus D$ has a neighbor in D . The domination number of G is the minimum cardinality of a dominating set of G . For a comprehensive survey of domination in graphs, see [2,3].

An edge $e \in E(G)$ is vertex-edge dominated by a vertex $v \in V(G)$ if e is incident to v , or e is adjacent to an edge incident to v . A subset $D \subseteq V(G)$ is a vertex-edge dominating set, abbreviated VEDS, of G if every edge of G is vertex-edge dominated by a vertex of D . The vertex-edge domination number of G , denoted by $\gamma_{ve}(G)$, is the minimum cardinality of a vertex-edge dominating set of G . A vertex-edge dominating set of G of minimum cardinality is called a $\gamma_{ve}(G)$ -set. Vertex-edge domination in graphs was introduced in [7], and further studied in [6].

Chellali and Haynes [1] established the following lower bound on the total domination number of a tree. For every tree T of order n with l leaves, we have $\gamma_t(T) \geq (n - l + 2)/2$. They also characterized the extremal trees. In [4] a lower bound on the total outer-independent domination number of a tree was given together with the characterization of the extremal trees. Lemańska [5] proved that the domination number of a tree is bounded below by $(n - l + 2)/3$.

We prove the following bounds on the vertex-edge domination number of a tree T of order $n \geq 3$ with l leaves and s support vertices, $(n - l - s + 3)/4 \leq \gamma_{ve}(T) \leq n/3$. We also characterize the trees attaining each of the bounds.

2. Results

We begin with the following straightforward observation.

Observation 1. For every connected graph G of diameter at least two, there exists a $\gamma_{ve}(G)$ -set that contains no leaf.

First we show that if T is a nontrivial tree of order n with l leaves and s support vertices, then $\gamma_{ve}(T)$ is bounded below by $(n - l - s + 3)/4$. For the purpose of characterizing the trees attaining this bound, we introduce a family \mathcal{T} of trees $T = T_k$ that can be obtained as follows. Let T_1 be a path P_5 . If k is a positive integer, then T_{k+1} can be obtained recursively from T_k by one of the following operations.

- Operation \mathcal{O}_1 : Attach a vertex by joining it to any support vertex of T_k .
- Operation \mathcal{O}_2 : Attach a path P_2 by joining one of its vertices to a vertex of T_k , which is not a leaf and is adjacent to a support vertex of degree two.
- Operation \mathcal{O}_3 : Attach a path P_4 by joining one of its leaves to a leaf of T_k adjacent to a weak support vertex.

We now prove that for every tree T of the family \mathcal{T} we have $\gamma_{ve}(T) = (n - l - s + 3)/4$.

Lemma 2. If $T \in \mathcal{T}$, then $\gamma_{ve}(T) = (n - l - s + 3)/4$.

Proof. We use the induction on the number k of operations performed to construct the tree T . If $T = T_1 = P_5$, then $(n - l - s + 3)/4 = (5 - 2 - 2 + 3)/4 = 1 = \gamma_{ve}(T)$. Let k be a positive integer. Assume that the result is true for every tree $T' = T_k$ of the family \mathcal{T} constructed by $k - 1$ operations. Let n' be the order of the tree T' , l' the number of its leaves, and s' the number of support vertices. Let $T = T_{k+1}$ be a tree of the family \mathcal{T} constructed by k operations.

First assume that T is obtained from T' by operation \mathcal{O}_1 . We have $n = n' + 1$, $l = l' + 1$ and $s = s'$. It is straightforward to see that any $\gamma_{ve}(T')$ -set is a VEDS of the tree T . Thus $\gamma_{ve}(T) \leq \gamma_{ve}(T')$. Obviously, $\gamma_{ve}(T') \leq \gamma_{ve}(T)$. This implies that $\gamma_{ve}(T) = \gamma_{ve}(T')$. We now get $\gamma_{ve}(T) = \gamma_{ve}(T') = (n' - l' - s' + 3)/4 = (n - 1 - l + 1 - s + 3)/4 = (n - l - s + 3)/4$.

Now assume that T is obtained from T' by operation \mathcal{O}_2 . We have $n = n' + 2$, $l = l' + 1$ and $s = s' + 1$. It is straightforward to see that any $\gamma_{ve}(T')$ -set is a VEDS of the tree T . Thus $\gamma_{ve}(T) \leq \gamma_{ve}(T')$. This implies that $\gamma_{ve}(T) = \gamma_{ve}(T')$. We now get $\gamma_{ve}(T) = \gamma_{ve}(T') = (n' - l' - s' + 3)/4 = (n - 2 - l + 1 - s + 1 + 3)/4 = (n - l - s + 3)/4$.

Now assume that T is obtained from T' by operation \mathcal{O}_3 . We have $n = n' + 4$, $l = l'$ and $s = s'$. We denote by x the leaf to which P_4 is attached. Let $v_1 v_2 v_3 v_4$ be the attached path. Let v_1 be joined to x . Let D' be any $\gamma_{ve}(T')$ -set. It is easy to see that $D' \cup \{v_2\}$ is a VEDS of the tree T . Thus $\gamma_{ve}(T) \leq \gamma_{ve}(T') + 1$. Now let us observe that there exists a $\gamma_{ve}(T)$ -set that does not contain the vertices v_4 , v_3 , and v_1 . Let D be such a set. To dominate the edge $v_3 v_4$, we have $v_2 \in D$. Observe that $D \setminus \{v_2\}$ is a VEDS of the tree T' . Therefore $\gamma_{ve}(T') \leq \gamma_{ve}(T) - 1$. We now conclude that $\gamma_{ve}(T) = \gamma_{ve}(T') + 1$. We get $\gamma_{ve}(T) = \gamma_{ve}(T') + 1 = (n' - l' - s' + 3)/4 + 1 = (n - 4 - l - s + 7)/4 = (n - l - s + 3)/4$. \square

We now give a lower bound on the vertex-edge domination number of a tree together with the characterization of the extremal trees.

Theorem 3. If T is a nontrivial tree of order n with l leaves and s support vertices, then $\gamma_{ve}(T) \geq (n - l - s + 3)/4$ with equality if and only if $T \in \mathcal{T}$.

Proof. If $\text{diam}(T) = 1$, then $T = P_2$. We have $(n - l - s + 3)/4 = (2 - 2 - 2 + 3)/4 < 1 = \gamma_{ve}(T)$. If $\text{diam}(T) = 2$, then T is a star. We have $l = n - 1$ and $s = 1$. Consequently, $(n - l - s + 3)/4 = (n - n + 1 - 1 + 3)/4 = 3/4 < 1 = \gamma_{ve}(T)$.

Download English Version:

<https://daneshyari.com/en/article/4669895>

Download Persian Version:

<https://daneshyari.com/article/4669895>

[Daneshyari.com](https://daneshyari.com)