



Mathematical Analysis/Numerical Analysis

Sharp inequalities related to Gosper's formula

Inégalités précises liées à la formule de Gosper

Cristinel Mortici

Valahia University of Târgoviște, Department of Mathematics, Bd. Unirii 18, 130082 Târgoviște, Romania

ARTICLE INFO

Article history:

Received 29 April 2009

Accepted after revision 10 December 2009

Available online 19 February 2010

Presented by Jean-Pierre Kahane

ABSTRACT

The purpose of this Note is to construct a new type of Stirling series, which extends the Gosper's formula for big factorials. New sharp inequalities for the gamma and digamma functions are established. Finally, numerical computations which demonstrate the superiority of our new series over the classical Stirling's series are given.

© 2010 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Le but de cette Note est de construire un nouveau type de série de Stirling, étendant la formule de Gosper pour les grandes factorielles. Nous établissons de nouvelles inégalités précises pour les fonctions gamma et digamma. Enfin, nous indiquons des calculs numériques qui démontre la supériorité de notre nouvelle série sur la série classique de Stirling.

© 2010 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

Probably one from the most known approximation formula for the factorial function is the Stirling formula:

$$n! \approx \sqrt{2\pi n} \cdot \left(\frac{n}{e}\right)^n,$$

that is the first estimate of the Stirling series, e.g., [1, p. 257],

$$n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \left(1 + \frac{1}{12n} + \frac{1}{288n^2} - \frac{139}{51840n^3} - \frac{517}{2488320n^4} + \dots\right). \quad (1)$$

In this Note we discuss the following more accurate formula:

$$n! \approx \sqrt{2\pi \left(n + \frac{1}{6}\right)} \cdot \left(\frac{n}{e}\right)^n \quad (\text{Gosper [4]}),$$

proving the following double inequality for $x \geq 1$,

$$\sqrt{2\pi \left(x + \frac{1}{6}\right)} \left(\frac{x}{e}\right)^x < \Gamma(x+1) \leq \omega \cdot \sqrt{2\pi \left(x + \frac{1}{6}\right)} \left(\frac{x}{e}\right)^x, \quad (2)$$

E-mail address: cmortici@valahia.ro.

where the constant $\omega = e\sqrt{\frac{3}{7\pi}} = 1.0039940821\dots$ is the best possible. Furthermore, we prove new sharp bounds for the digamma function ψ , e.g. [1, p. 258]. For $x \geq 1$, it holds

$$\ln x - \frac{1}{x} + \frac{1}{2(x + \frac{1}{6})} - \zeta \leq \psi(x) < \ln x - \frac{1}{x} + \frac{1}{2(x + \frac{1}{6})}, \quad (3)$$

with the best possible constant $\zeta = \gamma - \frac{4}{7} = 0.0057864\dots$ ($\gamma = 0.577215\dots$ is the Euler–Mascheroni constant), improving other known results [2,5–7,14] as

$$\ln x - \frac{1}{x} < \psi(x) < \ln x - \frac{1}{2x}, \quad x > 1.$$

Instead of completing the approximation series at the end part of the formula, as in the classical Stirling's series, we continue the series under the square root, the first two terms being given by the Gosper's formula. In this way, we obtain a series converging unexpectedly faster than the well-known Stirling series, namely

$$n! \approx \left(\frac{n}{e}\right)^n \sqrt{2\pi \left(n + \frac{1}{6} + \frac{1}{72n} - \frac{31}{6480n^2} - \frac{139}{155520n^3} + \frac{9871}{6531840n^4} + \dots\right)},$$

which can be viewed as an extension of the Stirling and Gosper formulas.

2. The results

In the first part of this section, we prove the following:

Theorem 1. The function $f : [1, \infty) \rightarrow \mathbb{R}$ given by

$$f(x) = \ln \Gamma(x+1) - x \ln x + x - \ln \sqrt{2\pi} - \frac{1}{2} \ln \left(x + \frac{1}{6}\right)$$

is convex, and strictly decreasing.

Proof. We have

$$f'(x) = \psi(x) + \frac{1}{x} - \ln x - \frac{1}{2(x + \frac{1}{6})}, \quad f''(x) = \psi'(x) - \frac{1}{x^2} - \frac{1}{x} + \frac{1}{2(x + \frac{1}{6})^2}.$$

According to a result of Gordon [3, Theorem 4], for every $x > 0$, we have

$$\psi'(x) > \frac{1}{x} + \frac{1}{2x^2} + \frac{1}{6x^3} - \frac{1}{30x^5}.$$

As for every $x \geq 1$,

$$\left(\frac{1}{x} + \frac{1}{2x^2} + \frac{1}{6x^3} - \frac{1}{30x^5}\right) - \left(\frac{1}{x^2} + \frac{1}{x} - \frac{1}{2(x + \frac{1}{6})^2}\right) = \frac{45 - \frac{31}{x} - \frac{12}{x^2} - \frac{1}{x^3}}{30x^8(6x+1)^2} > 0,$$

it results that $f''(x) > 0$, for every $x \geq 1$. Now, f' is strictly increasing with $\lim_{x \rightarrow \infty} f'(x) = 0$, thus $f' < 0$ and f is strictly decreasing. \square

As a direct consequence of the fact that f is strictly decreasing, we have $0 = \lim_{x \rightarrow \infty} f(x) < f(x) \leq f(1) = 1 + \ln \sqrt{\frac{3}{7\pi}}$, for every $x \geq 1$, which is the sharp inequality (2).

Using the fact that f' is strictly increasing, we have $\frac{4}{7} - \gamma = f'(1) \leq f'(x) < \lim_{x \rightarrow \infty} f'(x) = 0$, for every $x \geq 1$, which is the sharp inequality (3).

Now we are interested in finding the real values a, b, c, d that provide the most accurate formula of the form

$$n! \approx \left(\frac{n}{e}\right)^n \sqrt{2\pi \left(n + \frac{1}{6} + \frac{a}{n} + \frac{b}{n^2} + \frac{c}{n^3} + \frac{d}{n^4}\right)}. \quad (4)$$

To this end, we introduce the sequence $(\lambda_n)_{n \geq 1}$ by the relations

$$n! = \left(\frac{n}{e}\right)^n \sqrt{2\pi \left(n + \frac{1}{6} + \frac{a}{n} + \frac{b}{n^2} + \frac{c}{n^3} + \frac{d}{n^4}\right)} \cdot \exp \lambda_n, \quad (5)$$

and we say that an approximation (5) is better, the faster $(\lambda_n)_{n \geq 1}$ converges to zero. A basic tool for measuring the rate of convergence is the following:

Download English Version:

<https://daneshyari.com/en/article/4670988>

Download Persian Version:

<https://daneshyari.com/article/4670988>

[Daneshyari.com](https://daneshyari.com)