

Available online at www.sciencedirect.com

SciVerse ScienceDirect

**Procedia** Earth and Planetary Science

Procedia Earth and Planetary Science 7 (2013) 219 - 222

### Water Rock Interaction [WRI 14]

# Influence of water on rock discontinuities and stability of rock mass

Sandra Dochez<sup>1\*</sup>, Farid Laouafa<sup>1</sup>, Christian Franck<sup>2</sup> Sylvine Guedon<sup>3</sup>, François Martineau<sup>3</sup>, Marion Bost<sup>3</sup>, Julie D'Amato<sup>3</sup>

<sup>1</sup> INERIS, Parc Technologique Alata BP 2, Verneuil en Halatte 60550, France <sup>2</sup> GEODERIS, 1 rue Claude Chappe entrée C2, BP 25198, 57075 Metz Cedex 3, France. <sup>3</sup> IFSTTAR, Boulevard Newton, Champs sur Marne F-77447 Marne la Vallée Cedex 2, France

#### Abstract

Water is an important weathering factor that can play a role with respect to rock discontinuities, and thus rock mass behavior. The increase of rainfall (frequency and intensity) highlights new problems concerning rock mass stability. The aim of this research is an analysis of the importance of the role of water on rock alteration. This study will discuss how water causes discontinuity degradation, which can affect the stability of the rock mass. The accurate study of water-rock alteration is based upon two aspects, on the one hand the study of the intrinsic properties of water, and on the other hand, the study of the response of the discontinuity to alteration. The present study is based on two spatial scales, laboratory and in situ (the latter will not be discussed in this article).

© 2013 The Authors. Published by Elsevier B.V. Open access under CC BY-NC-ND license. Selection and/or peer-review under responsibility of the Organizing and Scientific Committee of WRI 14 – 2013

Keywords : weathering; chemical;, water; limestones; rock discontinuities; rock stability; roughness.

#### 1. Introduction

The stability of the rock mass depends on the presence of discontinuities, more specifically on their mechanical properties which are related to their morphological features such as opening, roughness. The study aims at evaluating the influence of water flow on the rock mass' stability. Water can be defined by many parameters; temperature, pH, salinity, hardness (bi-carbonate concentration) and many others [1]. We will observe the influence of the pH in two cases, when water is in a static mode or in dynamic mode. In the experiment where water is in a dynamic mode, we will observe the evolution of the roughness [5].

<sup>\*</sup> Corresponding author. Tel.: +33 3 44556283.

E-mail address: sandra.dochez@ineris.fr.

#### 2. Influence of pH on water alteration

A study carried out by Kupper (1985) proposed a relation between the rate of limestone erosion, the water flow velocity and the pH of the fluid. An equation was deduced [2]:

$$Log (Ve) = 5.103 + 0.423 (log (Vw)) - 0.442 pH$$

Where Ve is the erosion velocity expressed in [10<sup>-6</sup> m/years] and Vw is the water flow velocity expressed in [cm/s], pH is the pH of the solution (water).

In our experiments we choose an acidic solution at pH equal to 2 (sulfuric acid) whereas in situ pH is those of rain water around 5. This choice had been done in order to accelerate the chemical processes in laboratory. Using Kupper relation we obtain:

$$Ve(2) = 21 * Ve(5).$$

So the erosion velocity is multiplied by 21.

The pH is directly related to the concentration of [H+] and depends on the temperature.

We thought about two possibilities for alteration by water. Two experiments were made. In the first one, limestone samples are immerged in a solution of acidic water at room temperature. In the other, the solution of acidic water circulates on the surface's limestone samples at room temperature.





Figure 1: (a) Experiment 1 : limestone samples are drilled; drill-core is 10 cm long, and radius is 0.6 cm ; it is soaking in acidic water solution (b) Experiment 2: limestone samples are 5\*5\*8 cm parallelepiped with a rock bridge, of 1.5 cm of side; acidic water is added drop wise with a velocity of 1 liter per day.

The time evolution of the pH was measured during each of these experiments (Figure 2). We showed that the pH evolves differently. The experiment 1, showed an increase of the pH (2 to 7) whereas the pH remains around pH 2 during the experiment 2.

Download English Version:

## https://daneshyari.com/en/article/4675211

Download Persian Version:

https://daneshyari.com/article/4675211

Daneshyari.com