



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

Computers and Mathematics with Applications 55 (2008) 2999–3002

An International Journal  
**computers &  
mathematics**  
with applications

[www.elsevier.com/locate/camwa](http://www.elsevier.com/locate/camwa)

## Erratum

# Erratum to “Mann and Ishikawa iterative processes for multivalued mappings in Banach spaces” [Comput. Math. Appl. 54 (2007) 872–877]

Yisheng Song\*, Hongjun Wang

*College of Mathematics and Information Science, Henan Normal University, 453007, PR China*

Received 5 October 2007; accepted 17 November 2007

## Abstract

We show strong convergence for Mann and Ishikawa iterates of multivalued nonexpansive mapping  $T$  under some appropriate conditions, which revises a gap in Panyanak [B. Panyanak, Mann and Ishikawa iterative processes for multivalued mappings in Banach spaces, Comput. Math. Appl. 54 (2007) 872–877]. Furthermore, we also give an affirmative answer to Panyanak’s open question.

© 2008 Elsevier Ltd. All rights reserved.

**Keywords:** Ishikawa iterates; Strong convergence; Uniformly convex Banach spaces

Let  $E$  be a Banach space and  $K$  a nonempty subset of  $E$ . We shall denote  $CB(E)$  by the family of nonempty closed and bounded subsets of  $E$  and the family of nonempty bounded proximal subsets of  $E$  (see [1]). Let  $H$  be the Hausdorff metric on  $CB(E)$ , that is,

$$H(A, B) = \max\{\sup_{x \in A} d(x, B), \sup_{x \in B} d(x, A)\} \quad \text{for any } A, B \in CB(E),$$

where  $d(x, B) = \inf\{\|x - y\|; y \in B\}$ . A multivalued mapping  $T : K \rightarrow CB(E)$  is said to be *nonexpansive*, if for any  $x, y \in K$ , such that  $H(Tx, Ty) \leq \|x - y\|$ . A point  $x$  is called a fixed point of  $T$  if  $x \in Tx$ . From now on,  $F(T)$  stands for the fixed point set of a mapping  $T$ .

Recently, Panyanak [1] introduced the following Ishikawa iterates of a multivalued mapping  $T$ . Let  $K$  be a nonempty convex subset of  $E$ , fix  $p \in F(T)$  and  $x_0 \in K$ ,

$$y_n = (1 - \beta_n)x_n + \beta_n z_n, \quad \beta_n \in [0, 1], n \geq 0,$$

where  $z_n \in Tx_n$  such that  $\|z_n - p\| = d(p, Tx_n)$ , and

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n z'_n, \quad \alpha_n \in [0, 1], n \geq 0,$$

DOI of original article: [10.1016/j.camwa.2007.03.012](https://doi.org/10.1016/j.camwa.2007.03.012).

\* Corresponding author.

E-mail address: [songyisheng123@yahoo.com.cn](mailto:songyisheng123@yahoo.com.cn) (Y. Song).

where  $z'_n \in Ty_n$  such that  $\|z'_n - p\| = d(p, Ty_n)$ . It is obvious that  $x_n$  depends on  $p$  and  $T$ . For  $p \in F(T)$ , we have

$$\|z_n - p\| = d(p, Tx_n) \leq H(Tp, Tx_n) \leq \|x_n - p\|$$

and

$$\|z'_n - p\| = d(p, Ty_n) \leq H(Tp, Ty_n) \leq \|y_n - p\|.$$

Clearly, if  $q \in F(T)$  and  $q \neq p$ , then the above inequalities cannot be assured. Namely, from the monotony of  $\{\|x_n - p\|\}$  in the proof of [1, Theorem 3.1], we cannot obtain  $\{\|x_n - q\|\}$  is a decreasing sequence. Hence, the conclusion of Theorem 3.1 in [1] cannot be reached.

Motivated by solving the above gap, we have tried to modify it. The aim of this paper is to find an iteration instead of the above one and to overcome its limitation. We will construct the following iteration.

Let  $K$  be a nonempty convex subset of  $E$ ,  $\beta_n \in [0, 1]$ ,  $\alpha_n \in [0, 1]$  and  $\gamma_n \in (0, +\infty)$  such that  $\lim_{n \rightarrow \infty} \gamma_n = 0$ . Choose  $x_0 \in K$  and  $z_0 \in Tx_0$ . Let

$$y_0 = (1 - \beta_0)x_0 + \beta_0z_0.$$

There exists  $z'_0 \in Ty_0$  such that  $\|z_0 - z'_0\| \leq H(Tx_0, Ty_0) + \gamma_0$  (see [2,3]). Let

$$x_1 = (1 - \alpha_0)x_0 + \alpha_0z'_0.$$

There is  $z_1 \in Tx_1$  such that  $\|z_1 - z'_0\| \leq H(Tx_1, Ty_0) + \gamma_1$ . Take

$$y_1 = (1 - \beta_1)x_1 + \beta_1z_1.$$

There exists  $z'_1 \in Ty_1$  such that  $\|z_1 - z'_1\| \leq H(Tx_1, Ty_1) + \gamma_1$ . Let

$$x_2 = (1 - \alpha_1)x_1 + \alpha_1z'_1.$$

Inductively, we have

$$\begin{aligned} y_n &= (1 - \beta_n)x_n + \beta_nz_n, \\ x_{n+1} &= (1 - \alpha_n)x_n + \alpha_nz'_n, \end{aligned} \tag{1}$$

where  $\|z_n - z'_n\| \leq H(Tx_n, Ty_n) + \gamma_n$  and  $\|z_{n+1} - z'_n\| \leq H(Tx_{n+1}, Ty_n) + \gamma_n$  for  $z_n \in Tx_n$  and  $z'_n \in Ty_n$ .

We now show the strong convergence of the Ishikawa iteration (1) which shakes off the objection in [1, Theorem 3.1].

**Theorem 1.** Let  $K$  be a nonempty compact convex subset of a uniformly convex Banach space  $E$ . Suppose that  $T : K \rightarrow CB(K)$  is a multivalued nonexpansive mapping and  $F(T) \neq \emptyset$  satisfying  $T(y) = \{y\}$  for any fixed point  $y \in F(T)$ .

Let  $\{x_n\}$  be the sequence of Ishikawa iterates defined by (1). Assume that

(i)  $\alpha_n, \beta_n \in [0, 1]$ ; (ii)  $\lim_{n \rightarrow \infty} \beta_n = 0$  and (iii)  $\sum_{n=0}^{\infty} \alpha_n \beta_n = \infty$ .

Then as  $n \rightarrow \infty$ , the sequence  $\{x_n\}$  strongly converges to some fixed point of  $T$ .

**Proof.** Take  $p \in F(T)$  (noting  $Tp = \{p\}$  and  $\|z_n - p\| = d(z_n, Tp)$ ). Using a similar proof of Theorem 3.1 as in [1] (Xu's inequality, see [1, Lemma 2.3]), we have

$$\begin{aligned} \|x_{n+1} - p\|^2 &\leq (1 - \alpha_n)\|x_n - p\|^2 + \alpha_n\|z'_n - p\|^2 - \alpha_n(1 - \alpha_n)\varphi(\|x_n - z'_n\|) \\ &\leq (1 - \alpha_n)\|x_n - p\|^2 + \alpha_n(H(Ty_n, Tp))^2 \\ &\leq (1 - \alpha_n)\|x_n - p\|^2 + \alpha_n\|y_n - p\|^2 \\ &\leq (1 - \alpha_n)\|x_n - p\|^2 + \alpha_n[(1 - \beta_n)\|x_n - p\|^2 + \beta_n\|z_n - p\|^2 - \beta_n(1 - \beta_n)\varphi(\|x_n - z_n\|)] \\ &\leq (1 - \alpha_n)\|x_n - p\|^2 + \alpha_n[(1 - \beta_n)\|x_n - p\|^2 \\ &\quad + \beta_n(H(Tx_n, Tp))^2 - \beta_n(1 - \beta_n)\varphi(\|x_n - z_n\|)] \\ &\leq \|x_n - p\|^2 - \alpha_n\beta_n(1 - \beta_n)\varphi(\|x_n - z_n\|). \end{aligned}$$

Download English Version:

<https://daneshyari.com/en/article/470010>

Download Persian Version:

<https://daneshyari.com/article/470010>

[Daneshyari.com](https://daneshyari.com)