



Variational iteration method for twelfth-order boundary-value problems

A.S.V. Ravi Kanth*, K. Aruna

Applied Mathematics Division, School of Science and Humanities, V.I.T. University, Vellore – 632 014, Tamil Nadu, India

ARTICLE INFO

Keywords:

Variational iteration method
Linear–nonlinear problems
Boundary-value problems
Integral equations

ABSTRACT

In this paper, He's variational iteration method is applied to solve twelfth-order boundary-value problems. The numerical results obtained with minimum amount of computation are compared with the exact solutions to show the efficiency of the method. The results show that the variational iteration method is of high accuracy, more convenient and efficient for solving high-order boundary-value problems.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

We consider the twelfth-order boundary-value problem of the type:

$$y^{xii}(x) + f(x)y = g(x) \quad a < x < b \quad (1)$$

$$y^{2n}(0) = \alpha_{2n}, \quad y^{2n}(1) = \beta_{2n}, \quad n = 0, 1, \dots, 5 \quad (2)$$

where α_{2n} and β_{2n} ($n = 0, 1, \dots, 5$) are finite real constants and the functions $f(x)$ and $g(x)$ are continuous on $[a, b]$. This type of boundary-value problem arises in an infinite horizontal layer of fluid is heated from below, with the assumption that a uniform magnetic field is also applied across the fluid in the same direction as gravity and the fluid is subject to the action of rotation, instability sets in. When instability sets in as ordinary convection, it is modeled by tenth-order boundary-value problems, when instability sets in as over stability, it is modeled by twelfth-order boundary-value problems (see Chandrasekhar [1]). Agarwal's book [2] contains theorems which detail the conditions for existence and uniqueness of solutions of the twelfth-order boundary-value problems. Different numerical methods have been proposed by various authors for twelfth-order boundary-value problems. To mention a few, finite difference method [3,4], modified decomposition method [5], twelfth degree splines [6] and thirteen degree splines [7].

In this paper, we aim to apply the variational iteration method proposed by He [8–14] to twelfth-order boundary-value problems. This method has been employed to solve a large variety of linear and nonlinear problems such as the KdV, the K(2, 2), the Burgers and the cubic Boussinesq equations [23], Korteweg–de vries (KdV) and the modified Korteweg–de vries equations [16], nonlinear differential equations of fractional order [17], KdV Burgers equations and Lax's seventh-order KdV equations [18], generalized Burgers–Fisher and Burgers equations [19], Helmholtz equation [20], semi-linear inverse parabolic equation [15], linear fractional differential equation [21], nonlinear partial differential equations [22] etc.

2. Variational iteration method

According to the variational iteration method [8–14], we can construct a correction functional for twelfth-order boundary-value problem (1)–(2) in the form

$$y_{n+1}(x) = y_n(x) + \int_0^x \lambda [y_n^{xii}(s) + f(s)y_n(s) - g(s)] ds \quad (3)$$

* Corresponding author.

E-mail address: asvrvikanth@yahoo.com (A.S.V. Ravi Kanth).

where λ is a general Lagrange multiplier which can be identify via variational theory [8–10]. Making $y_{n+1}(x)$ stationary with respect to $y_n(x)$, we can identify the Lagrange multiplier, which reads

$$\lambda = \frac{(s-x)^{11}}{11!}. \quad (4)$$

Example 1. First, we consider the following BVP [7]

$$y^{xii}(x) + xy(x) + (120 + 23x + x^3)e^x = 0 \quad 0 < x < 1 \quad (5)$$

with boundary conditions

$$\left. \begin{aligned} y(0) = 0 = y(1), y'(0) = 1, y'(1) = -e, y''(0) = 0, \\ y''(1) = -4e, y'''(0) = -3, y'''(1) = -9e, y^{iv}(0) = -8, \\ y^{iv}(1) = -16e, y^v(0) = -15, y^v(1) = -25e \end{aligned} \right\}. \quad (6)$$

According to (3), we have the following iteration formulation

$$y_{n+1}(x) = y_n(x) + \int_0^x \frac{(s-x)^{11}}{11!} [y_n^{xii}(s) + sy_n(s) + (120 + 23s + s^3)e^s] ds. \quad (7)$$

Now, we assume that an initial approximation has the form

$$y_0(x) = a + bx + cx^2 + dx^3 + ex^4 + fx^5 + gx^6 + hx^7 + ix^8 + jx^9 + kx^{10} + lx^{11} \quad (8)$$

where $a, b, c, d, e, f, g, h, i, j, k$ and l are unknown constants to be further determined. By iteration formula (7), we have the following first-order approximation

$$\begin{aligned} y_1(x) &= y_0(x) + \int_0^x \frac{(s-x)^{11}}{11!} [y_0^{xii}(s) + sy_0(s) + (120 + 23s + s^3)e^s] ds \\ &= -\frac{l}{1295295050649600}x^{24} - \frac{k}{647647525324800}x^{23} - \frac{j}{309744468633600}x^{22} \\ &\quad - \frac{i}{140792940288000}x^{21} - \frac{h}{60339831552000}x^{20} - \frac{g}{24135932620800}x^{19} \\ &\quad - \frac{f}{8892185702400}x^{18} - \frac{e}{2964061900800}x^{17} - \frac{d}{871782912000}x^{16} - \frac{c}{217945728000}x^{15} \\ &\quad - \frac{b}{43589145600}x^{14} - \frac{a}{6227020800}x^{13} + \left(l + \frac{13}{5702400}\right)x^{11} + \left(k + \frac{1}{72576}\right)x^{10} \\ &\quad + \left(j - \frac{1}{40320}\right)x^9 + \left(i - \frac{23}{10080}\right)x^8 + \left(h - \frac{41}{1008}\right)x^7 + \left(g - \frac{59}{120}\right)x^6 + \left(f - \frac{109}{24}\right)x^5 \\ &\quad + \left(e - \frac{98}{3}\right)x^4 + \left(d - \frac{359}{2}\right)x^3 + (c - 715)x^2 + (b - 1849)x + (a - 2340) \\ &\quad + (2340 - 491x + 36x^2 - x^3)e^x. \end{aligned}$$

Incorporating the boundary conditions, Eq. (6), into $y_1(x)$, and solving the system simultaneously, we obtain

$$\begin{aligned} a &= 0, & b &= 1, & c &= 0, & d &= -0.5, & e &= -1/3, & f &= -1/8, & g &= -0.03333333362911, \\ h &= -0.00694444319430, & i &= -0.00119047835708, & j &= -0.00017360919905, \\ k &= -0.00002204671105, & l &= -0.00000248000384. \end{aligned}$$

We, therefore obtain the following first-order approximate solution

$$\begin{aligned} y_1(x) &= \frac{0.0000024899384}{1295295050649600}x^{24} + \frac{0.00002204671105}{647647525324800}x^{23} \\ &\quad + \frac{0.00017360919905}{309744468633600}x^{22} + \frac{0.001190478357105}{140792940288000}x^{21} + \frac{0.00694444319430}{60339831552000}x^{20} \\ &\quad + \frac{0.03333333362911}{24135932620800}x^{19} + \frac{1/8}{8892185702400}x^{18} + \frac{1/3}{2964061900800}x^{17} \\ &\quad + \frac{1/2}{87178291200}x^{16} - \frac{1}{43589145600}x^{14} - 2.00261976924803e-007x^{11} \end{aligned}$$

Download English Version:

<https://daneshyari.com/en/article/471909>

Download Persian Version:

<https://daneshyari.com/article/471909>

[Daneshyari.com](https://daneshyari.com)