




Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com



Original article

A new genus of male cones of voltzialean affinity, *Uralostrobus voltzioides* nov. gen., nov. sp., from the Lower Permian of the Urals (Russia)[☆]



Serge V. Naugolnykh^{a,b,*}

^a Geological Institute of Russian Academy of Sciences, 119017 Pyzhevsky per. 7, Moscow, Russia

^b Kazan Federal University, 18, Kremlyovskaya St., Kazan 420008, Republic of Tatarstan, Russian Federation

ARTICLE INFO

Article history:

Received 16 December 2013

Accepted 17 July 2014

Available online 2 October 2014

Keywords:

In situ pollen

New genus and species

Voltziaceae

Voltziales

Conifers

Gymnosperms

ABSTRACT

A new genus and species of male cones of coniferophyte from the Lower Permian (Artinskian and Kungurian) deposits of the Urals, Russia is described: *Uralostrobus voltzioides* Naugolnykh nov. gen., nov. sp. The cone shows characters typical of some representatives of conifers belonging to the order Voltziales: more or less isometrical bracts of rhombic shape, prolonged sporangia with attenuate apices, and bisaccate pollen of *Illinites*-type. General information on the associated female seed scales and vegetative leafy shoots is given as well.

© 2014 Elsevier Masson SAS. All rights reserved.

1. Introduction

Coniferophytes as a group of higher plants were widespread in the Permian period, especially in the Northern hemisphere, and after that they began to be intensively and rapidly diversified, which resulted in the appearance of several families, typical of the Triassic but having their evolutionary roots deep in the Permian (Clement-Westerhof, 1974; Miller, 1982; Meyen, 1987, 1997; Kerp et al., 1990; Mapes and Rothwell, 1991; Looy et al., 1999; Rothwell et al., 2005; Hernandez-Castillo et al., 2009). Traditionally, in palaeobotanical studies of Palaeozoic conifers the main attention is given to their female reproductive organs. Compared to the female cones and their isolated dwarf shoots of Permian conifers, during recent decades only a few publications were focused directly and exclusively on the male cones of that plant group. Nonetheless, the male reproductive organs are significant for taxonomy and systematics of early conifers, especially if information about their *in situ* pollen is obtained.

The present paper focuses on the description of coniferophyte male cones, which were found in the Lower Permian deposits of the Urals, Perm and Sverdlovsk regions (Russia). The *in situ* pollen grains

were extracted from the best preserved cone, which was selected as a holotype. All the male cones described here are attributed to a new genus and species, *Uralostrobus voltzioides* Naugolnykh nov. gen., nov. sp., most probably of voltzialean affinity.

2. Material and methods

The material studied includes four well-preserved specimens of male cones originating from three Early Permian localities from the Middle and Southern Cis-Urals (Fig. 1; Table 1; regional geology is considered after Chuvashov et al., 1990), in the Perm and Sverdlovsk Regions, Russia:

- Locality A: **Sobolevsky quarry**. This locality is disposed near the City of Krasnoufmsk (Sverdlovsk region), 3 km southwest from the Krasnoufmsk railway station. Stratigraphically, it belongs to the Lower Permian, Artinskian stage, Upper Artinskian Substage, Sargian Horizon (= Sargian Regional Stage), Divjinskian Fm. (= Divjinskaya Suite);
- Locality C: **Rakhmangulovo**. This locality is a small quarry near Rakhmangulovo Village, Krasnoufmsk District, 20 km east of the City of Krasnoufmsk. It belongs to the Lower Permian, Kungurian Stage, Lower Kungurian Substage, Philippovian Horizon (= Philippovian Regional Stage), Sabanakovian Fm. (= Sabanakovskaya Suite);

[☆] Corresponding editor: Marc Philippe.

* Geological Institute of Russian Academy of Sciences, 119017 Pyzhevsky per. 7, Moscow, Russia.

E-mail addresses: naugolnykh@rambler.ru, naugolnykh@list.ru.




Fig. 1. Geographical position of the localities studied and stratigraphical columns of the most important outcrops (Geologic map adapted from [Muzafarov, 1979](#)). Artinskian localities: A: Sobolevsky quarry; B: Shaidury; Kungurian localities: C: Rakhmangulovo; D: Krasnaya Glinka; E: Chekarda-1; F: Aleksandrovskoe. Legend for lithology: 1: sandstone; 2: aleurolite; 3: argillite; 4: clayey marl; 5: marl; 6: discontinuity with pebbles; 7: lens-like concretions; 8: cephalopods; 9: brachiopods; 10: terrestrial plants; 11: geographical position of the localities studied. Scale for stratigraphical columns: 1 m.

- Locality E: **Chekarda-1**. This locality is in the middle stream of the Sylva River (Suksun District, Perm region), 70 m downstream from the mouth of the Chekarda-River. Stratigraphically, it belongs to the Lower Permian, Kungurian Stage, Upper Kungurian Substage, Irenian Horizon (= Irenian Regional Stage), Koshelevskian Fm. (= Koshelevskaya Suite).

Other localities cited in the present paper are:

- Locality B: **Shaidury** (stems and leafy shoots probably of voltzialean affinity). Midstream of the river Sylva, Sverdlovsk region, Shalinsky district, 3 km upstream from the village

Download English Version:

<https://daneshyari.com/en/article/4748154>

Download Persian Version:

<https://daneshyari.com/article/4748154>

[Daneshyari.com](https://daneshyari.com)