


## Accepted Manuscript

Low-level laser therapy (LLLT) accelerates the sternomastoid muscle regeneration process after myonecrosis due to bupivacaine

Cristiane Neves Alessi Pissulin, Ana Angélica Henrique Fernandes, Alejandro Manuel Sanchez Orellana, Renata Calciolari Rossi e Silva, Selma Maria Michelin Matheus



PII: S1011-1344(16)30773-4  
DOI: doi: [10.1016/j.jphotobiol.2017.01.021](https://doi.org/10.1016/j.jphotobiol.2017.01.021)  
Reference: JPB 10722

To appear in: *Journal of Photochemistry & Photobiology, B: Biology*

Received date: 8 September 2016

Accepted date: 19 January 2017

Please cite this article as: Cristiane Neves Alessi Pissulin, Ana Angélica Henrique Fernandes, Alejandro Manuel Sanchez Orellana, Renata Calciolari Rossi e Silva, Selma Maria Michelin Matheus , Low-level laser therapy (LLLT) accelerates the sternomastoid muscle regeneration process after myonecrosis due to bupivacaine. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Jpb*(2017), doi: [10.1016/j.jphotobiol.2017.01.021](https://doi.org/10.1016/j.jphotobiol.2017.01.021)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**LOW-LEVEL LASER THERAPY (LLLT) ACCELERATES THE  
STERNOMASTOID MUSCLE REGENERATION PROCESS AFTER  
MYONECROSIS DUE TO BUPIVACAINE**

**Cristiane Neves Alessi Pissulin M.D.<sup>1</sup>, Ana Angélica Henrique Fernandes M.D,PhD<sup>2</sup>, Alejandro Manuel Sanchez Orellana<sup>3</sup>, Renata Calciolari Rossi e Silva PhD.<sup>4</sup>, Selma Maria Michelin Matheus M.D., PhD.<sup>5</sup>**

<sup>1</sup>Department of Anatomy, Universidade do Oeste Paulista (UNOESTE), Presidente Prudente, SP, Brazil; General Bases of Surgery, Botucatu Medical School; Unesp, Botucatu, SP, Brasil, crispissulin@gmail.com.

<sup>2</sup>Department of Chemistry and Biochemistry, Institute of Bioscience, Unesp, Botucatu, SP, Brasil; angelica@ibb.unesp.br.

<sup>3</sup>Biological Sciences Student, Unesp, Institute of Bioscience, Botucatu, SP, Brasil, alesanc96@outlook.com.

<sup>4</sup>Department of Pathology, Universidade do Oeste Paulista (UNOESTE), Presidente Prudente, SP, Brasil, renata@unoeste.br.

<sup>5</sup>Department of Anatomy, Institute of Bioscience; General Bases of Surgery, Botucatu Medical School; Unesp, Botucatu, SP, Brasil, micmath@ibb.unesp.br.

**Correspondence**

Address correspondence to Dra. Cristiane Neves Alessi Pissulin, Departamento de Anatomia, Faculdade de Odontologia. Rua José Bongiovanni, 700, Cep 18050-680, Unoeste, Campus I/ Presidente Prudente /SP/Brasil, crispissulin@gmail.com; +55(018)32291092.

**Keywords:** Bupivacaine, Low-Level Light Therapy, Muscle Regeneration, Fibrosis.

**Highlights:**

- . LLLT is a co-adjuvant therapy after lesions triggered by Bupivacaine.
- . LLLT accelerates the process of muscle regeneration.
- . LLLT reduce fibrosis and myonecrosis (sternomastoid muscle) caused by bupivacaine.

**Competing Interests**

The authors declare no competing interests.

**Research Support:** This work was supported by Fapesp Grant n° 13/26649-3

**Acknowledgment:** PhD. Carlos Roberto Padovani for Statistical analyses.

Download English Version:

<https://daneshyari.com/en/article/4754530>

Download Persian Version:

<https://daneshyari.com/article/4754530>

[Daneshyari.com](https://daneshyari.com)