


Accepted Manuscript

Selective oxidation of 2-naphthol to 2-hydroxy-1,4-naphthoquinone with hydrogen peroxide catalyzed by 5,10,15,20-tetrakis(p-sulfonato-phenyl)porphinat manganese(III) chloride in aqueous solution.



M. Hassanein, S. El-Khalafy, S. Shendy

PII: S1566-7367(17)30102-4

DOI: doi: [10.1016/j.catcom.2017.03.016](https://doi.org/10.1016/j.catcom.2017.03.016)

Reference: CATCOM 4973

To appear in: *Catalysis Communications*

Received date: 11 December 2016

Revised date: 16 March 2017

Accepted date: 18 March 2017

Please cite this article as: M. Hassanein, S. El-Khalafy, S. Shendy , Selective oxidation of 2-naphthol to 2-hydroxy-1,4-naphthoquinone with hydrogen peroxide catalyzed by 5,10,15,20-tetrakis(p-sulfonato-phenyl)porphinat manganese(III) chloride in aqueous solution.. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Catcom(2017), doi: [10.1016/j.catcom.2017.03.016](https://doi.org/10.1016/j.catcom.2017.03.016)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Selective oxidation of 2-naphthol to 2-hydroxy-1,4-naphthoquinone with hydrogen peroxide catalyzed by 5,10,15, 20-tetrakis(p- sulfonato-phenyl)porphinat manganese(III) chloride in aqueous solution.

M. Hassanein*, S. El-Khalafy, S. Shendy

Tanta University, Faculty of Science, Department of Chemistry, Tanta 31527, Egypt.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/4756633>

Download Persian Version:

<https://daneshyari.com/article/4756633>

[Daneshyari.com](https://daneshyari.com)