



New records of forensic entomofauna in legally buried and exhumed human infants remains in Buenos Aires, Argentina



Roxana Mariani ^{a,*}, Rocío García-Mancuso ^b, Graciela L. Varela ^a, Ivana Kierbel ^a

^a División Entomología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque, 1900 La Plata, Buenos Aires, Argentina

^b Cátedra de Citología, Histología y Embriología "A", Facultad de Ciencias Médicas, Universidad Nacional de La Plata, Calle 60 y 120, 1900 La Plata, Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 29 May 2017

Received in revised form

15 September 2017

Accepted 18 September 2017

Available online 22 September 2017

Keywords:

Forensic entomology

Underground cadaver colonization

Forensic anthropology

Legal graves

ABSTRACT

The study of carrion fauna associated with buried human corpses from a forensic perspective could provide useful information in criminal investigations. Insects and other arthropods remains sampled of 44 legally exhumed infant skeletons from La Plata (Buenos Aires, Argentina). They were identified at different taxonomic levels depending on the state of preservation. The specific diversity, abundance and frequency were analyzed and each taxon was assigned to the hypothetical colonization sequence: burial colonization, post-exhumation contamination at cemetery deposit or soil fauna. The phorid *Dohrniphora* sp. is mentioned for the first time in Argentina as carrion fauna of underground colonization, and the assemblage of *Dohrniphora* sp., *Megaselia scalaris* and *Hydrotaea aenescens* is proposed as indicator of buried cadavers. These findings provide new useful data to be applied in forensic entomology research.

© 2017 Elsevier Ltd and Faculty of Forensic and Legal Medicine. All rights reserved.

1. Introduction

Forensic entomologists collaborate in legal investigations; it studies the insects and other arthropods found in decomposing bodies and uses them to determine the moment and circumstances of death.¹ This discipline considers that the corpse (from death to the skeletal stage) functions as a true ecosystem that is colonized in a predictable pattern, known as ecological succession, which can offer relevant information in criminal proceedings and trials.^{2,3}

The colonization process has been studied mainly on above-ground outdoor and indoor animal carcasses^{2,4–6} but also on human bodies.^{7–11} The patterns of this faunal colonization may vary according to cadaver characteristics, such as size, weight,¹² the presence of wounds, or environmental conditions as sun exposure, season of the year and geographical region, which also have an effect on arrival times and tenure of different insects' species.^{13–15}

Studies of insect colonization on buried bodies has been carried out using carcasses at different depths^{16–18} whilst other came from exhumations of human corpses buried in soil or in coffins.^{19–22} Burial depth and the type of soil could influence in the

accessibility to the body, and the underground lower temperatures and oxygen availability produce a slower decomposition rate of bodies and reduced dispersion of decay odours; all of these could delayed the colonization and reduced the insects specific diversity that could reach the remains.^{16,20,23–25} Recent research carried out in field and laboratory experiments analyzed the effect of burial on the development of larvae and pupae of forensically important flies in different types of soils and confirmed the ability of phorids,²⁶ muscids and calliphorids²⁷ and sarcophagids²⁸ to colonize buried remains.

In Argentina, the first mention of necrophagous entomofauna of legal exhumations from crypts and soil was made by Oliva,²¹ who mentioned *Hydrotaea argentina* (Bigot, 1885), actually synonymy of *H. aenescens* (Wiedemann, 1830), *Muscina* sp. and *Megaselia scalaris* (Loew, 1866) for the city of Buenos Aires. Recently, Mariani et al.²² provided an inventory of entomofauna recovered from the skeleton of a young child and the funerary bundle belonging to a legal exhumation from the ground in the cemetery of La Plata (Buenos Aires Province), in which the muscid *Hydrotaea aenescens* (cited as *Ophyra aenescens*), was the dominant taxon. They proposed a hypothetical colonization sequence, one during the burial in soil and then a post-exhumation contamination at the cemetery deposit.

The aim of the present study was to register the occurrences of entomofauna associated to legal exhumations of human infant skeletons from Argentina and assess the taxa that can be used as

* Corresponding author.

E-mail addresses: rmariani@fcnym.unlp.edu.ar (R. Mariani), rgarciamancuso@gmail.com (R. García-Mancuso), gracielaavarela02@gmail.com (G.L. Varela), ivanakierbel@hotmail.com (I. Kierbel).

indicators of buried cadavers in forensic cases and criminal investigations.

2. Materials and methods

The faunal remains analyzed in this paper were sampled from 44 exhumed skeletonized bodies of fetuses and infants up to one year old. The bodies were buried shortly after death and from individual death records from the cemetery archives it is known that inhumations happened all around the year and the bodies kept buried between 4 and 8 years. After this period of time, exhumation took place and the skeletons were stored at the cemetery deposit for relatives or the administration to decide upon their destination (Ordenanza Municipal 7638/90). After a period of time of between 2 and 7 years in the cemetery deposit, those skeletons that went unclaimed were donated by the Municipal Cemetery of La Plata (Cementerio Municipal de La Plata) to the School of Medical Sciences (Facultad de Ciencias Médicas, FCM), National University of La Plata (Universidad Nacional de La Plata, UNLP) for research and teaching purposes (Ordenanza Municipal 9471/02). The skeletons arrived at FCM in numbered plastic bags including clothing, external wrappings and sediments; they were cleaned and prepared for their inclusion in the 'Prof. Dr. Rómulo Lambre' skeletal collection housed in the FCM where they currently belong.²⁹

2.1. Burial conditions

The burial location in the city of La Plata, Buenos Aires Province (34° 57' 20" S, 57° 57' 5.96" W) has the humid temperate climate of the Pampas region, a mean annual precipitation of 1146 mm, and a marked seasonality. The soil is silty and, at a depth of 50 cm it has an annual average temperature of 17.2 °C (11 °C in winter and 21.4 °C in summer).³⁰ The inhumation occurred in the four seasons of the year and bodies were buried in the ground, sometimes clothed, placed in a soft wooden coffin (specially made for rapid decomposition) in a 40-cm-deep grave, and covered with loose soil.

2.2. Arthropod remains

During the process of conditioning of the skeletal material before its inclusion in the collection's storage, the fauna was sampled using brushes, pins or micro forceps, depending on the specimens' size and conditions, and the sediments were inspected through water flotation. These consisted of whole or fragmented insect adults, pupae, exuviae of larvae and other arthropods, and molluscan shells. The samples obtained were stored in labeled sealed plastic bags.

Morphological and systematic studies were undertaken at the Entomology Division (División de Entomología), School and Museum of Natural Sciences (Facultad de Ciencias Naturales y Museo, FCNyM, UNLP), where dry specimens were observed under a stereoscopic microscope (Leica EZ5). The identification was possible at the species, family or order taxonomic levels depending on the preservation of the material.

Forensically importance dipteran remains were identified by three authors (RM, GLV and IK). The characters used for the identification were based on the posterior segments, the relative position of the tubercles and features of the posterior spiracles; their shape and distance apart; the contours of the spiracular slits; the thickness, shape and completeness of the peritreme and the location of the scar and button. All these characters were diagnosed according to Queiroz and de Carvalho³¹; Lui and Greenberg³²; Oliva^{33,34}; Sukontason et al.³⁵; Barbosa Cortinhas et al.³⁶; Grzywacz et al.^{37, 38}; Feng and Liu.³⁹ Coleopterans were identified according to Almeida and Mise.⁴⁰ The exemplars were compared against

reference specimens of La Plata Museum collection (MLP, FCNyM, UNLP). Voucher specimens have been deposited in the collection of the MLP.

To facilitate the interpretation of the analysis of the data in the "Results" section, each exhumed skeleton was described as a "case". All of them were buried in soil inside a coffin in a particular unsheltered area of the cemetery destined to infants, and after exhumation they were placed in the cemetery deposit into numbered plastic bags that constitute their container until inclusion in Lambre collection.

The arrival, establishment and development of the entomofauna was interpreted on the basis of the location and depth of the burials, previous forensic records, and the biology of each taxon.²² A first underground colonization (burial colonization) characterized by carrion species capable of laying eggs on the remains surface and the neonated larvae reaching corpses by boring through the substrate. A second instance of colonization (post-exhumation colonization), dominated by omnivorous and storage pests that occurred in the context of the cemetery deposit and characterized by insects unable to burrow as adults or larvae. Finally, soil fauna of the burial substrate sampled with the remains is described.

The faunal attributes analyzed were: abundance, which was recorded as the number of specimens of a given taxon in the whole sample of human remains; frequency, considered as the number of occurrences of a taxon per skeleton. The association of cadaver fauna analyzed per skeleton and found in more than one case was mentioned as assemblage.

3. Results

A total of 1216 specimens were collected from the 44 cases. Carrion fauna known to colonize buried corpses were the most numerous (n = 1025) and were registered in 28 cases. Other specimens were attributed to contamination at the cemetery deposit or accidental arrivals (n = 49), occurring in 20 cases, and in 24 cases the typical soil fauna of the burial substrate (n = 142) was found. Finally, cadaver fauna was found together with the fauna of the deposit in nine cases and with soil fauna in fifteen cases. The inventory of faunal remains is presented in Table 1.

3.1. Burial insect colonization

As regards cadaver fauna, the most abundant and frequent taxon was *Megaselia scalaris* (Fig. 1a), followed in importance by other fly remains of *Hydrotaea aenescens* (Fig. 1b), *Dohrniphora* sp. (Fig. 1c), and less abundant *Muscina stabulans* (Fallén, 1817) (Fig. 1d), and the coleopteran *Ataenius* sp. (Fig. 2). *M. scalaris*, *H. aenescens* and *Dohrniphora* sp. were found together and with a high number of specimens and also, *M. scalaris* and *H. aenescens* were often found together or along with less frequent taxa (*M. stabulans* and *Ataenius* sp. respectively) (Fig. 3). Some puparia were observed within other puparia; specifically, *M. scalaris* and *H. aenescens* puparia were found within puparia of their same species. Furthermore, in three cases, puparia of the above-mentioned species were also found inside bone cavities (Fig. 1e).

3.2. Post-exhumation insect colonization

At the cemetery deposit, the moth *Tineola bisselliella* (Hummel, 1823) was the most abundant and frequent insect; *Tinea pellionella* (Linnaeus, 1758) was the other lepidopteran recorded. Among beetles, the most abundant was *Anobium punctatum* (De Geer, 1774) followed by *Anthrenus verbasci* (Linnaeus, 1767), and *Dermestes* sp. In addition, cockroach oothecae (Blattodea), and remains of earwig (Anisolabididae) were found (Table 1).

Download English Version:

<https://daneshyari.com/en/article/4760619>

Download Persian Version:

<https://daneshyari.com/article/4760619>

[Daneshyari.com](https://daneshyari.com)