

Contents lists available at ScienceDirect

Data in Brief





Data Article

Genotoxicity assessment data for exfoliated buccal cells exposed to mobile phone radiation



F.M. de Oliveira *, A.M. Carmona, C. Ladeira

Institute of Cancer Research, School of Tourism and Maritime Technology, Lisbon School of Health Technology, United Kingdom

ARTICLE INFO

Article history:
Received 5 August 2017
Received in revised form
8 September 2017
Accepted 20 September 2017
Available online 22 September 2017

Keywords: Electromagnetic fields Mobile phones Genotoxicity Micronuclei Exfoliated buccal cells Feulgen stain

ABSTRACT

Healthy mobile phone users aged 18–30 y.o. provided exfoliated buccal cells samples from the right and left inner cheeks. A total of 2000 cells per subject were screened for the presence of micronuclei as a sign of genotoxic damage, according to the mobile phone use profile of each user.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

Specifications Table

Subject area Physics, Biology
More specific subject Radiation genotoxicity

area

Type of data Text file, graphs

How data was Leica DM500 Microscope, survey

acquired

Data format Analyzed

Experimental factors Histochemical stain with Feulgen's method

DOI of original article: http://dx.doi.org/10.1016/j.mrgentox.2017.08.001

 $^{\ast}\ Corresponding\ author.$

E-mail address: flavia.mendesdeoliveira@icr.ac.uk (F.M. de Oliveira).

Experimental features

Data source location
Data accessibility
Related research article

Aircroscope screening at 1000x with immersion oil

Lisbon Metropolitan Area, Portugal, 38.7223 ° N, 9.1393 ° W

Data are within this article.

Genotoxicity assessment of mobile phone radiation in exfoliated buccal cells (in press)

Value of the data

- Data was collected from members of an important share of mobile phone users (young adults, aged 18–30). Establishing the effects of mobile phone use in this population can contribute to an overall perception of how such devices affect the majority of its users [1].
- Improved perception of the effects of mobile phone electromagnetic radiation in humans can contribute to improved safety guidelines for the use of this device and help combat long standing misconceptions on mobile phone radiation [2–5].
- Establishing the relevance and efficacy of exposure levels and of the biomarker assessment method
 herein described can help in the determination of a genotoxicity-based model of observation and
 thus promote the development of new methods.

1. Data

Overall micronucleus frequency in the study population (2.02 (\pm 1.65) per 2000 cells) was found to be within currently accepted physiological ranges [6]. Lifestyle factors assessed in subjects were not shown to affect the frequency of this genotoxicity biomarker, with the exception of occupational exposure to known genotoxic agents (Fig. 1). Daily duration, side of use and history of mobile phone in years (Fig. 2) did not correlate to higher micronucleus frequencies.

2. Experimental design, materials and methods

Buccal exfoliated cells were collected using sterile endobrushes followed by a smearing technique on histological slides. Cells were fixated with an ethanol-based solution, air-dried and stained according to Feulgen's method [7]. Mounted slides were screened by a singleobserver at a 1000x magnification with immersion oil and morphological objects within accepted intervals for




Fig. 1. Density plot of micronuclei frequency in subjects exposed and non-exposed to known genotoxic agents.

Download English Version:

https://daneshyari.com/en/article/4764905

Download Persian Version:

https://daneshyari.com/article/4764905

<u>Daneshyari.com</u>