



Data Article

Data for differentially expressed microRNAs in saturated fatty acid palmitate-treated HepG2 cells



Won-Mo Yang^a, Kyung-Ho Min^a, Wan Lee^{a,b,*}

^a Department of Biochemistry, Dongguk University College of Medicine, Gyeongju 780-714, South Korea

^b Endocrine Channelopathy, Channelopathy Research Center, Dongguk University College of Medicine, Goyang, 410-773, South Korea

ARTICLE INFO

Article history:

Received 10 October 2016

Received in revised form

16 November 2016

Accepted 16 November 2016

Available online 19 November 2016

Keywords:

MicroRNAs

Palmitate

Saturated fatty acids

Obesity

Hepatocyte

ABSTRACT

Certain microRNAs (miRNAs) targeting the molecules in the insulin signaling cascades are dysregulated by saturated fatty acids (SFA), which can lead to insulin resistance and type 2 diabetes. This article reports the accompanying data collected using miRNAs microarrays to identify the changes in miRNA expression in HepG2 cells treated with SFA palmitate. Differentially expressed miRNA analyses in HepG2 cells showed that a range of upregulated (> 1.5 -fold) or downregulated (< 0.5 -fold) miRNAs. Further extensive insights into the implications of miRNAs, particularly miR-1271, in HepG2 cells can be found in "MiR-1271 upregulated by saturated fatty acid palmitate provokes impaired insulin signaling by repressing INSR and IRS-1 expression in HepG2 cells" (W.M. Yang, K.H. Min, W. Lee, 2016) [1].

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

| | |
|----------------------------|-------------------------------|
| Subject area | Biology, Biochemistry |
| More specific subject area | Obesity, Metabolism, MicroRNA |

DOI of original article: <http://dx.doi.org/10.1016/j.bbrc.2016.09.029>

* Corresponding author at: Department of Biochemistry, Dongguk University College of Medicine, Gyeongju 780-714, Korea.

E-mail address: wanlee@dongguk.ac.kr (W. Lee).

<http://dx.doi.org/10.1016/j.dib.2016.11.062>

2352-3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

| | |
|-----------------------|--|
| Type of data | <i>Excel files</i> |
| How data was acquired | <i>Affymetrix GeneChip microarray analyses of miRNAs</i> |
| Data format | <i>Analyzed</i> |
| Experimental factors | <i>Palmitate treatment, RNA Isolation, Affymetrix Genechip miRNA microarray</i> |
| Experimental features | <i>Differentially expressed miRNAs of the HepG2 cells treated with SFA palmitate were analyzed using Affymetrix GeneChip miRNA microarray.</i> |
| Data source location | <i>Dongguk University School of Medicine, Gyeongju 780-714, Korea</i> |
| Data accessibility | <i>The data are available with this article</i> |

Value of the data

- The data highlight the biological significance of the miRNAs involved in the pathogenesis of SFA-induced metabolic diseases.
 - These results can be compared with gene expression analysis from other cell or tissue types in obesity.
 - The differentially expressed miRNAs in this dataset could be applied in further functional studies of the cellular and systemic phenotype changes resulting from SFA-induced obesity and metabolic diseases.
-

1. Data

The high dietary intake of saturated fatty acids (SFA), which is the leading cause of obesity, frequently causes ectopic lipid accumulation and increase the risk of insulin resistance in non-adipose tissues, such as the liver and skeletal muscle [2]. The expression of certain miRNAs targeting the insulin signaling molecules is modulated aberrantly in diet-induced obesity, which participates actively in the pathogenesis of insulin resistance [3,4]. A previous study reported that SFA palmitate induces miR-1271 in HepG2 hepatocytes, and the expression of INSR and IRS-1 is suppressed by targeting their 3'UTR directly [1]. This means that certain miRNA induced by SFA could be linked causally to the development of hepatic insulin resistance and further to type 2 diabetes. This paper reports accompanying data collected from Affymetrix GeneChip microarrays to identify the changes in miRNA expression in HepG2 cells treated with SFA palmitate. Differentially expressed microRNA analyses in HepG2 cells (Supplementary File. 1) revealed a range of miRNAs upregulated more than 1.5-fold (Supplementary File. 2) or downregulated less than 0.5-fold (Supplementary File. 3). Among those differentially expressed miRNAs, upregulated miRNAs have implications on the reduction of INSR and IRS-1 observed in palmitate-treated HepG2 cells [1]. Further analysis of the data and insights into the implications of miRNAs, especially miR-1271, in HepG2 cells are reported in another publication [1].

2. Experimental design, materials and methods

2.1. Cells and palmitate treatment

HepG2, a human liver cancer cell line, was purchased from ATCC (#77400). The HepG2 cells were grown in MEM α supplemented with 10% FBS and 1% penicillin-streptomycin (Gibco) in an atmosphere containing 5% CO₂ at 37 °C. The cells from passages 3–10 were used for the following experiments. A fatty acid-free bovine serum albumin (BSA, Bovogen, VIC, Australia)-conjugated palmitate (Sigma-Aldrich) solution was prepared, as described previously [5]. Briefly, BSA and sodium palmitate were dissolved completely in 150 mM NaCl by heating at 37 °C and 70 °C, respectively. The BSA solution was added dropwise to the palmitate solution at 37 °C with continuous stirring until the

Download English Version:

<https://daneshyari.com/en/article/4765451>

Download Persian Version:

<https://daneshyari.com/article/4765451>

[Daneshyari.com](https://daneshyari.com)