

Accepted Manuscript

Fluorescent anticancer quinazolines as molecular probes for β -tubulin colchicine site competition assay and visualization of microtubules as intracellular targeting sites

Yumiko Suzuki, Jun-ichi Sawada, Paulina Hibner, Hirosuke Ishii, Kenji Matsuno, Masayuki Sato, Bernhard Witulski, Akira Asai



PII: S0143-7208(17)30461-8

DOI: [10.1016/j.dyepig.2017.05.050](https://doi.org/10.1016/j.dyepig.2017.05.050)

Reference: DYPI 6012

To appear in: *Dyes and Pigments*

Received Date: 14 March 2017


Revised Date: 1 May 2017

Accepted Date: 29 May 2017

Please cite this article as: Suzuki Y, Sawada J-i, Hibner P, Ishii H, Matsuno K, Sato M, Witulski B, Asai A, Fluorescent anticancer quinazolines as molecular probes for β -tubulin colchicine site competition assay and visualization of microtubules as intracellular targeting sites, *Dyes and Pigments* (2017), doi: 10.1016/j.dyepig.2017.05.050.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The distinguished fluorophore performance of quinazoline **1** in non-polar hydrophobic media allows its use as molecular probe for β -tubulin site competition assays and for visualization of its intracellular distribution.



Download English Version:

<https://daneshyari.com/en/article/4765776>

Download Persian Version:

<https://daneshyari.com/article/4765776>

[Daneshyari.com](https://daneshyari.com)