

Separation of binary mixtures hexane/hex-1-ene, cyclohexane/cyclohexene and ethylbenzene/styrene based on limiting activity coefficients

Urszula Domańska, Michal Wlazło, Monika Karpińska, Maciej Zawadzki

PII: S0021-9614(17)30064-2
DOI: http://dx.doi.org/10.1016/j.jct.2017.03.004
Reference: YJCHT 4998

To appear in: *J. Chem. Thermodynamics*

Received Date: 27 January 2017
Revised Date: 6 March 2017
Accepted Date: 8 March 2017

Please cite this article as: U. Domańska, M. Wlazło, M. Karpińska, M. Zawadzki, Separation of binary mixtures hexane/hex-1-ene, cyclohexane/cyclohexene and ethylbenzene/styrene based on limiting activity coefficients, *J. Chem. Thermodynamics* (2017), doi: http://dx.doi.org/10.1016/j.jct.2017.03.004

J. Chem. Thermodyn.

Separation of binary mixtures hexane/hex-1-ene, cyclohexane/cyclohexene and ethylbenzene/styrene based on limiting activity coefficients

Urszula Domańska[a,b]*, Michal Wlazło[c], Monika Karpińska[c], Maciej Zawadzki[c]

[a]*Industrial Chemistry Research Institute, Rydygiera 8, 01-793 Warsaw, Poland.*

[b] *Thermodynamic Research Unit, School of Chemical Engineering, University of KwaZulu-Natal, Howard College Campus, King George V Avenue, Durban 4001, South Africa.*c

[c] *Department of Physical Chemistry, Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warsaw, Poland*

Keywords:

Ionic liquid, [*N*-C_3OHPY][DCA]
Experimental limiting activity coefficients

Thermodynamics

Separation

The Abraham solvation parameter model

*Corresponding author at Department of Physical Chemistry, Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warsaw, Poland. Tel.:+48 22 6213115; fax: +48 22 6282741. *E-mail address*: ula@ch.pw.edu.pl (U. Domańska).