



Cairo University
Egyptian Informatics Journal

www.elsevier.com/locate/eij
www.sciencedirect.com



ORIGINAL ARTICLE

Rational trigonometric cubic spline to conserve convexity of 2D data

Farheen Ibraheem ^{a,*}, Maria Hussain ^b, Malik Zawwar Hussain ^c

^a National University of Computer and Emerging Sciences, Lahore, Pakistan

^b Lahore College for Women University, Lahore, Pakistan

^c Department of Mathematics, University of the Punjab, Lahore, Pakistan

Received 25 March 2013; revised 12 June 2013; accepted 7 July 2013

Available online 5 August 2013

KEYWORDS

Rational trigonometric cubic function;
Free parameters;
Convex data;
Data visualization

Abstract Researchers in different fields of study are always in dire need of spline interpolating function that conserve intrinsic trend of the data. In this paper, a rational trigonometric cubic spline with four free parameters has been used to retain convexity of 2D data. For this purpose, constraints on two of free parameters β_i and γ_i in the description of the rational trigonometric function are derived while the remaining two α_i and δ_i are set free. Numerical examples demonstrate that resulting curves using the technique of the underlying paper are C^1 .

© 2013 Production and hosting by Elsevier B.V. on behalf of Faculty of Computers and Information, Cairo University.

1. Introduction

Representation of data in the form of congenial curves and surfaces is of great importance in many areas of scientific research such as computer graphics and data visualization. This visual display of data provides prompt cognition and insight into data. Shape preservation and smoothness are the most desirable features required by a researcher in the field of data

visualization. Spline interpolating function demonstrates the incredible result in this regard.

Properties that quantify shape of data are positivity, monotonicity, and convexity. Plenty of spline functions exist which can produce a smooth curve but inept to preserve the inherent shape of the given data. The motivation of this paper is to preserve the intrinsic attribute of data that is convexity.

Convexity pervades everyday life. Whether it is manufacturing of lenses, modeling of cars, analysis of indifference curves, nonlinear programming and approximation of functions, convexity stays the part and parcel of all. Loss of convexity is irreconcilable in all such practical problems.

Over the years, many milestones have been achieved in the field of shape preservation when data under consideration exhibit convex trend. Various schemes have been developed to reach the epitome of abstraction. Brodlie and Butt [1] preserved the convex shape of data by establishing a piecewise cubic function. Their scheme suffered the detriment of insertion of the additional knots in an interval where convexity is lost.

* Corresponding author. Tel.: +92 42 99231444.

E-mail addresses: farheen.butt@gmail.com (F. Ibraheem), mariahus-sain_1@yahoo.com (M. Hussain), malikzawwar.math@pu.edu.pk (M.Z. Hussain).

Peer review under responsibility of Faculty of Computers and Information, Cairo University.



Production and hosting by Elsevier

Convexity preserving properties of rational Bezier and non-uniform rational B-spline were studied geometrically by Carnicer [2]. Explicit representation of rational cubic function with one free parameter was developed by Delbourgo and Gregory [3] to retain convexity. Passcow and Roulier [4] formed a spline interpolant by constructing an auxiliary set of points and using convexity preserving properties of Bernstein polynomials. Schumaker [5] preserved the convexity by piecewise quadratic polynomial which was economical but again an extra knot had to be inserted. A rational cubic function with two shape parameters was introduced by Sarfraz and Hussain [6] to retain convexity. Sarfraz et al. [7] developed rational cubic functions with four parameters. In [7], the authors derived the shape preserving constraints on two parameters and two parameters are left free for the user to refine the shape of the curves.

The algorithm presented in this paper is a noteworthy addition to already existing schemes and is a leap forward in many ways. The use of trigonometric function strengthens a designer to frame conics accurately. It does not require insertion of extra knots to conserve the intrinsic trend of the data. Four free parameters have been used in the specification of the rational trigonometric cubic spline. Constraints on two of free parameters are derived to envision convex data. The remaining two free parameters give enough freedom to the user for the refinement of convex shape of data according to the requirement.

The remainder of the paper is organized as follows. Section 2 is a review of the rational trigonometric cubic function [8] to be used for convexity. Section 3 deals with the development of convexity conserving constraints. In Section 4, numerical examples have been demonstrated. Section 5 summarizes the contributions and concludes the paper.

2. Rational trigonometric cubic function

Let $\{(x_i, f_i), i = 0, 1, 2, \dots, n\}$ be the given set of data points defined over the interval $[a, b]$ where $a = x_0 < x_1 < x_2 < \dots < x_n = b$. A piecewise rational trigonometric cubic function is defined over each subinterval $I_i = [x_i, x_{i+1}]$ as

$$S_i(x) = \frac{p_i(\theta)}{q_i(\theta)}, \quad (1)$$

$$\begin{aligned} p_i(\theta) = & \alpha_i f_i (1 - \sin \theta)^3 + \left\{ \beta_i f_i + \frac{2h_i \alpha_i d_i}{\pi} \right\} \sin \theta (1 - \sin \theta)^2 \\ & + \left\{ \gamma_i f_{i+1} - \frac{2h_i \delta_i d_{i+1}}{\pi} \right\} \cos \theta (1 - \cos \theta)^2 \\ & + \delta_i f_{i+1} (1 - \cos \theta)^3, \end{aligned}$$

$$\begin{aligned} q_i(\theta) = & \alpha_i (1 - \sin \theta)^3 + \beta_i \sin \theta (1 - \sin \theta)^2 + \gamma_i \cos \theta (1 - \cos \theta)^2 \\ & + \delta_i (1 - \cos \theta)^3, \end{aligned}$$

where

$$\theta = \frac{\pi}{2} \left(\frac{x - x_i}{h_i} \right), \quad h_i = x_{i+1} - x_i, \quad i = 0, 1, 2, \dots, n-1.$$

The rational trigonometric cubic function (1) satisfies the following properties:

$$S(x_i) = f_i, \quad S(x_{i+1}) = f_{i+1}, \quad S^{(1)}(x_i) = d_i, \quad S^{(1)}(x_{i+1}) = d_{i+1}.$$

d_i and d_{i+1} are derivatives at the end points of the interval $I_i = [x_i, x_{i+1}]$, α_i , β_i , γ_i and δ_i are the free parameters.

3. Convex rational trigonometric cubic spline

In this section, problem of preserving convexity of 2D is addressed. For this purpose, constraints on free parameters in the description of rational trigonometric cubic function (1) are developed. Let the convex data set defined over the interval $[a, b]$ be $\{(x_i, f_i), i = 0, 1, 2, \dots, n\}$. The necessary condition for the convexity of data is

$$d_1 < \Delta_1 < \dots < \Delta_{i-1} < d_i < \Delta_i < \dots < \Delta_{n-1} < d_n.$$

The rational trigonometric cubic function (1) is convex if and only if $S_i''(x) > 0$, where

$$\begin{aligned} S_i''(x) = & \frac{\pi}{2h_i(q_i(\theta))^2} \{ A_0 \sin \theta (1 - \sin \theta)^8 + A_1 \cos^2 \theta (1 - \sin \theta)^7 \\ & + A_2 \sin^2 \theta (1 - \sin \theta)^7 + A_3 \sin \theta \cos^2 \theta (1 - \sin \theta)^6 \\ & + A_4 \sin \theta \cos^2 \theta (1 - \sin \theta)^5 (1 - \cos \theta) \\ & + A_5 \sin^2 \theta \cos \theta (1 - \sin \theta)^6 + A_6 \sin \theta \cos^2 \theta (1 - \sin \theta)^6 \\ & + A_7 \sin^2 \theta (1 - \sin \theta)^6 (1 - \cos \theta) \\ & + A_8 \cos \theta (1 - \sin \theta)^6 (1 - \cos \theta)^2 \\ & + A_9 \sin^3 \theta \cos \theta (1 - \sin \theta)^5 \\ & + A_{10} \sin^2 \theta \cos^2 \theta (1 - \sin \theta)^4 (1 - \cos \theta) \\ & + A_{11} \sin^3 \theta (1 - \sin \theta)^5 (1 - \cos \theta) \\ & + A_{12} \sin^2 \theta \cos \theta (1 - \sin \theta)^4 (1 - \cos \theta)^2 \\ & + A_{13} \sin \theta \cos \theta (1 - \sin \theta)^5 (1 - \cos \theta)^2 \\ & + A_{14} \cos^3 \theta (1 - \sin \theta)^4 (1 - \cos \theta)^2 \\ & + A_{15} \cos^3 \theta \sin \theta (1 - \sin \theta)^3 (1 - \cos \theta)^2 \\ & + A_{16} \cos \theta \sin^2 \theta (1 - \sin \theta)^3 (1 - \cos \theta)^3 \\ & + A_{17} \cos^2 \theta (1 - \sin \theta)^3 (1 - \cos \theta)^4 \\ & + A_{18} \cos^2 \theta \sin \theta (1 - \sin \theta)^3 (1 - \cos \theta)^3 \\ & + A_{19} \sin^2 \theta (1 - \sin \theta)^3 (1 - \cos \theta)^4 \\ & + A_{20} \cos^2 \theta (1 - \sin \theta)^4 (1 - \cos \theta)^3 \\ & + A_{21} \sin \theta (1 - \sin \theta)^5 (1 - \cos \theta)^3 \\ & + A_{22} \sin^2 \theta (1 - \sin \theta)^4 (1 - \cos \theta)^3 \\ & + A_{23} \cos \theta (1 - \sin \theta)^3 (1 - \cos \theta)^5 \\ & + A_{24} \sin^3 \theta (1 - \sin \theta)^6 + A_{25} \sin^2 \theta \cos^2 \theta (1 - \sin \theta)^5 \\ & + A_{26} \sin^4 \theta \cos \theta (1 - \sin \theta)^4 \\ & + A_{27} \sin^3 \theta (1 - \cos \theta)^4 (1 - \sin \theta)^2 \\ & + A_{28} \sin \theta \cos \theta (1 - \cos \theta)^5 (1 - \sin \theta)^2 \\ & + A_{29} \sin^3 \theta \cos \theta (1 - \cos \theta)^3 (1 - \sin \theta)^2 \\ & + A_{30} \sin^3 \theta \cos \theta (1 - \cos \theta)^2 (1 - \sin \theta)^3 \\ & + A_{31} \sin^4 \theta (1 - \cos \theta) (1 - \sin \theta)^4 \\ & + A_{32} \sin^3 \theta \cos^2 \theta (1 - \cos \theta) (1 - \sin \theta)^3 \\ & + A_{33} \sin \theta \cos^2 \theta (1 - \cos \theta)^4 (1 - \sin \theta)^2 \\ & + A_{34} \sin^2 \theta \cos^3 \theta (1 - \cos \theta)^2 (1 - \sin \theta)^2 \\ & + A_{35} \sin^3 \theta (1 - \cos \theta)^3 (1 - \sin \theta)^3 \\ & + A_{36} \sin^2 \theta \cos^2 \theta (1 - \cos \theta)^3 (1 - \sin \theta)^2 \\ & + A_{37} \sin^2 \theta \cos \theta (1 - \cos \theta)^6 + A_{38} \cos^2 \theta (1 - \cos \theta)^7 \\ & + A_{39} \sin^2 \theta \cos^2 \theta (1 - \cos \theta)^4 (1 - \sin \theta) \\ & + A_{40} \cos^3 \theta (1 - \cos \theta)^3 (1 - \sin \theta)^3 \\ & + A_{41} \sin^2 \theta \cos^2 \theta (1 - \cos \theta)^2 (1 - \sin \theta)^3 \\ & + A_{42} \sin \theta \cos^3 \theta (1 - \cos \theta)^3 (1 - \sin \theta)^2 \\ & + A_{43} \sin^3 \theta \cos^2 \theta (1 - \cos \theta)^2 (1 - \sin \theta)^2 \\ & + A_{44} \cos^3 \theta (1 - \cos \theta)^6 + A_{45} \cos^4 \theta (1 - \cos \theta)^4 (1 - \sin \theta) \\ & + A_{46} \cos^3 \theta (1 - \cos \theta)^5 (1 - \sin \theta) \\ & + A_{47} \cos^4 \theta \sin \theta (1 - \cos \theta)^4 + A_{48} \sin^2 \theta \cos \theta (1 - \cos \theta)^5 (1 - \sin \theta) \\ & + A_{49} \sin \theta \cos^3 \theta (1 - \cos \theta)^5 + A_{50} \sin^2 \theta (1 - \cos \theta)^3 (1 - \sin \theta)^4 \\ & + A_{51} \cos \theta (1 - \cos \theta)^8 + A_{52} \sin^2 \theta (1 - \cos \theta)^7 \\ & + A_{53} \sin \theta (1 - \cos \theta)^3 (1 - \sin \theta)^5 + A_{54} \sin \theta (1 - \cos \theta)^6 (1 - \sin \theta)^2 \\ & + A_{55} \sin \theta \cos^2 \theta (1 - \cos \theta)^3 (1 - \sin \theta)^3 + A_{56} \cos^2 \theta (1 - \cos \theta)^6 (1 - \sin \theta) \\ & + A_{57} \sin^2 \theta (1 - \cos \theta)^6 (1 - \sin \theta) + A_{58} \sin \theta \cos^2 \theta (1 - \cos \theta)^6 \\ & + A_{59} \sin^2 \theta \cos^2 \theta (1 - \sin \theta)^5 + A_{60} \sin^2 \theta \cos^2 \theta (1 - \cos \theta)^3 (1 - \sin \theta)^2 \\ & + A_{61} \sin^3 \theta \cos^2 \theta (1 - \cos \theta) (1 - \sin \theta)^3 + A_{62} \sin^2 \theta \cos^3 \theta (1 - \cos \theta)^3 (1 - \sin \theta) \\ & + A_{63} \sin^2 \theta \cos^2 \theta (1 - \cos \theta)^4 (1 - \sin \theta) \} \end{aligned} \quad (2)$$

Download English Version:

<https://daneshyari.com/en/article/477357>

Download Persian Version:

<https://daneshyari.com/article/477357>

[Daneshyari.com](https://daneshyari.com)