



Egyptian Mathematical Society
Journal of the Egyptian Mathematical Society

www.etms-eg.org
www.elsevier.com/locate/joems



ORIGINAL ARTICLE

A new multivalued contraction and stability of its fixed point sets



Binayak S. Choudhury¹, Chaitali Bandyopadhyay *

Department of Mathematics, Bengal Engineering and Science University, Shibpur, Howrah 711103, India

Received 8 December 2013; revised 20 April 2014; accepted 6 May 2014

Available online 11 June 2014

KEYWORDS

Metric space;
Fixed point set;
Hausdorff metric;
Multivalued α - ψ
contraction;
Stability

Abstract In this paper we obtain some stability results for fixed point sets associated with a sequence of multivalued mappings. We define multivalued α - ψ contractions and multivalued α -admissible mappings. We use Hausdorff distance in our definition. We show that the fixed point sets of uniformly convergent sequences of multivalued α - ψ contractions which are also assumed to be multivalued α -admissible, are stable under certain conditions. The multivalued mappings we define here are not necessarily continuous. We present two illustrative examples and one open problem.

2010 MATHEMATICS SUBJECT CLASSIFICATION: 47H10; 37C75; 54C60

© 2014 Production and hosting by Elsevier B.V. on behalf of Egyptian Mathematical Society.

1. Introduction and preliminaries

Stability is a concept in dynamical systems related to limiting behaviors. There are various notions of stability both in discrete and continuous dynamical systems [1,2]. In this paper we consider such a problem of stability related to a sequence of multivalued mappings on metric spaces. The limiting behaviors of sequences of mappings have been considered in a large number of papers in recent times as, for instances, in

[3,4]. Particularly, stability of fixed point sets for multivalued mapping has been considered in [5–7].

Specially, we are interested in the limit of fixed point sets for a convergent sequence of multivalued mappings, that is, how they are related, in the limit, to the fixed point set of the function to which the sequence converges. We say that the fixed point sets are stable when they converge in the Hausdorff metric to the set of fixed points of the limiting function. More often than not, in the above mentioned problem of stability, sequences of multivalued mappings are considered. One of the reasons behind this is that multivalued mappings often have more fixed points than their singlevalued counterparts. For instance, in the theorem of Nadler [3], which is a multivalued generalization of the Banach contraction principle, and, incidentally, which is also the first work appearing on multivalued contractive fixed point studies, the fixed point is not unique in contrast to the case of Banach's contraction. In those situations the fixed point set becomes larger and, hence, more interesting for the study of stability. In this paper

* Corresponding author. Tel.: +91 9433471023.

E-mail addresses: binayak12@yahoo.co.in (B.S. Choudhury), chaitali.math@gmail.com (C. Bandyopadhyay).

¹ Tel.: +91 9433800840; fax: +91 33 26682916.

Peer review under responsibility of Egyptian Mathematical Society.



we consider the case of α - ψ contractions [8] which is a newly introduced generalization of the Banach's contraction. It should be mentioned that Banach contraction mapping principle [9,10] plays an important role in nonlinear analysis. There has been a large number of generalizations of this result over the years [11–16].

We introduce a multivalued version of α - ψ contraction. We show that for such a multivalued mapping on a complete metric space, the fixed point set is nonempty. We then show that a uniformly convergent sequence of such mappings on a complete metric space has stable fixed point sets, that is, the fixed point sets converge to the fixed point set of the limiting function with respect to the Hausdorff metric.

Throughout this paper $CL(X)$ denotes the family of all nonempty closed subsets of a metric space (X, d) and $P(X)$ denotes the family of all nonempty subsets of X .

The Hausdorff metric H is defined on $CL(X)$ by

$$H(A, B) = \max \left\{ \sup_{x \in B} d(x, A), \sup_{x \in A} d(x, B) \right\},$$

where $A, B \in CL(X)$ and $d(x, A) = \inf_{y \in A} d(x, y)$.

H is a metric when it is restricted to the set $CB(X)$, the set of all closed and bounded subsets of X . Otherwise, on $CL(X)$, the set of all closed subsets of X , all the properties of the metric function is satisfied except that $H(A, B)$ can be infinite when either A or B is unbounded. The following is the well known definition of fixed point for multivalued mappings.

Let $T : X \rightarrow P(X)$ be a multivalued mapping, a point $z \in X$ is a fixed point of T whenever $z \in Tz$.

Asl et al. [17] introduce the following definition;

Definition 1.1 [17]. Let (X, d) be a metric space; $\alpha : X \times X \rightarrow [0, \infty)$ be a mapping and $T : X \rightarrow 2^X$ be a closed valued multifunction, where $2^X =$ collection of all nonempty subsets of X . Let $\psi : [0, \infty) \rightarrow [0, \infty)$ be a nondecreasing and continuous function with $\sum \psi^n(t) < \infty$ and $\psi(t) < t$ for each $t > 0$. We say that T is an α - ψ contractive multifunction whenever

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \psi(d(x, y)), \text{ for } x, y \in X, \quad (1.1)$$

where $\alpha_*(Tx, Ty) = \inf\{\alpha(a, b) : a \in Tx, b \in Ty\}$.

In the following we introduce the concept of multivalued α - ψ contraction and multivalued α -admissible.

Definition 1.2 (Multivalued α - ψ contraction). Let (X, d) be a metric space, and $\alpha : X \times X \rightarrow [0, \infty)$, $\psi : [0, \infty) \rightarrow [0, \infty)$ be two mappings such that ψ is a nondecreasing and continuous function with $\sum \psi^n(t) < \infty$ and $\psi(t) < t$ for each $t > 0$. $T : X \rightarrow CL(X)$ be a multivalued mapping. We say that T is a multivalued α - ψ contraction if

$$\alpha(x, y)H(Tx, Ty) \leq \psi(d(x, y)), \text{ for all } x, y \in X. \quad (1.2)$$

Remark 1.1. In (1.2) of our Definition 1.2 we consider $\alpha(x, y)$ instead of $\alpha_*(Tx, Ty)$ which has been considered in (1.1) of Definition 1.1. $\alpha_*(Tx, Ty)$ is defined as

$$\alpha_*(Tx, Ty) = \inf\{\alpha(a, b) : a \in Tx, b \in Ty\}, \text{ for } x, y \in X.$$

From the definition it is clear that $\alpha_*(Tx, Ty)$ is not necessarily equal to $\alpha(x, y)$, and also we cannot compare $\alpha(x, y)$ with

$\alpha_*(Tx, Ty)$. Therefore Definition 1.2 is new and independent of Definition 1.1.

Remark 1.2. If T is singlevalued in Definition 1.2, then it is an α - ψ contraction as in [8].

Definition 1.3 (Multivalued α -admissible). Let X be any nonempty set. $T : X \rightarrow P(X)$ and $\alpha : X \times X \rightarrow [0, \infty)$ be two mappings. We say that T is multivalued α -admissible if, for $x, y \in X$,

$$\alpha(x, y) > 1 \Rightarrow \alpha(a, b) > 1, \text{ for all } a \in Tx \text{ and for all } b \in Ty.$$

Example 1.1. Let $X = \mathbb{R}$, $\alpha : \mathbb{R} \times \mathbb{R} \rightarrow [0, \infty)$. We define

$$\alpha(x, y) = x^2 + y^2, \text{ where } x, y \in \mathbb{R}.$$

Define $T : \mathbb{R} \rightarrow P(\mathbb{R})$ by,

$$Tx = \left\{ \sqrt{|x|}, -\sqrt{|x|} \right\}.$$

Then T is multivalued α -admissible.

2. Main Result

We first prove that multivalued α - ψ contractions on complete metric spaces have nonempty fixed point sets. In the proof of the following theorem we make use of Lemma 8.1.3(c) of [18].

Theorem 2.1. Let (X, d) be a complete metric space and $T : X \rightarrow CL(X)$ be a multivalued α - ψ contraction. Also T satisfies the following:

- (i) T is multivalued α -admissible;
- (ii) For some $x_0 \in X$, $\alpha(x_0, a) > 1$ holds for all $a \in Tx_0$;
- (iii) If $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) > 1$ for all n , where $x_{n+1} \in Tx_n$ and $x_n \rightarrow x$ as $n \rightarrow \infty$, then $\alpha(x_n, x) > 1$ for all n .

Then T has a fixed point.

Proof. Let $x_0 \in X$ be as in the statement of the theorem. By (ii), we have $x_1 \in Tx_0$ such that, $\alpha(x_0, x_1) > 1$. Then, since $x_1 \in Tx_0$, we can choose $x_2 \in Tx_1$ such that, $d(x_1, x_2) \leq \alpha(x_0, x_1)H(Tx_0, Tx_1)$. So, by (1.2), we have

$$d(x_1, x_2) \leq \alpha(x_0, x_1)H(Tx_0, Tx_1) \leq \psi(d(x_0, x_1)). \quad (2.1)$$

Since $x_1 \in Tx_0, x_2 \in Tx_1$ and $\alpha(x_0, x_1) > 1$, by (i), we have $\alpha(x_1, x_2) > 1$.

Again, for $x_2 \in Tx_1$, we can choose $x_3 \in Tx_2$ such that

$$d(x_2, x_3) \leq \alpha(x_1, x_2)H(Tx_1, Tx_2).$$

Therefore, by (1.2), we have,

$$\begin{aligned} d(x_2, x_3) &\leq \alpha(x_1, x_2)H(Tx_1, Tx_2) \leq \psi(d(x_1, x_2)) \\ &\leq \psi^2(d(x_0, x_1)) \quad (\text{by (2.1)}). \end{aligned} \quad (2.2)$$

Also, since $\alpha(x_1, x_2) > 1, x_2 \in Tx_1$ and $x_3 \in Tx_2$ we have that $\alpha(x_2, x_3) > 1$. Continuing this process we can construct

Download English Version:

<https://daneshyari.com/en/article/483446>

Download Persian Version:

<https://daneshyari.com/article/483446>

[Daneshyari.com](https://daneshyari.com)