

# Accepted Manuscript

Comparison of Vis or NIR Hyperspectral Imaging and Computer Vision for Automatic Differentiation of Organically and Conventionally Farmed Salmon

Jun-Li Xu, Cecilia Riccioli, Da-Wen Sun



PII: S0260-8774(16)30382-X  
DOI: [10.1016/j.jfoodeng.2016.10.021](https://doi.org/10.1016/j.jfoodeng.2016.10.021)  
Reference: JFOE 8698  
To appear in: *Journal of Food Engineering*  
Received Date: 24 August 2015  
Revised Date: 21 September 2016  
Accepted Date: 20 October 2016

Please cite this article as: Jun-Li Xu, Cecilia Riccioli, Da-Wen Sun, Comparison of Vis or NIR Hyperspectral Imaging and Computer Vision for Automatic Differentiation of Organically and Conventionally Farmed Salmon, *Journal of Food Engineering* (2016), doi: 10.1016/j.jfoodeng.2016.10.021

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

### Highlights

- CVS and Vis/NIR HSI were investigated to classify salmon fillets into four grades.
- Colour and spectral feature differences of samples were extracted and analysed.
- PLS-DA, SVM and RF classifiers were applied and compared in discriminant analysis.
- Optimal variables were selected by iPLSDA in visible and NIR region, respectively.
- SVM model for the 400-1000 nm HSI yielded best result for prediction.

Download English Version:

<https://daneshyari.com/en/article/4909212>

Download Persian Version:

<https://daneshyari.com/article/4909212>

[Daneshyari.com](https://daneshyari.com)