


Accepted Manuscript

Title: Exploring suicide risk factors among Japanese individuals: the largest case-control psychological autopsy study in Japan

Authors: Manami Kodaka, Toshihiko Matsumoto, Michiko Takai, Takashi Yamauchi, Shizuka Kawamoto, Minako Kikuchi, Hisateru Tachimori, Yotaro Katsumata, Norihito Shirakawa, Tadashi Takeshima



PII: S1876-2018(17)30004-7
DOI: <http://dx.doi.org/doi:10.1016/j.ajp.2017.02.029>
Reference: AJP 1079

To appear in:

Received date: 4-1-2017
Revised date: 23-2-2017
Accepted date: 23-2-2017

Please cite this article as: Kodaka, Manami, Matsumoto, Toshihiko, Takai, Michiko, Yamauchi, Takashi, Kawamoto, Shizuka, Kikuchi, Minako, Tachimori, Hisateru, Katsumata, Yotaro, Shirakawa, Norihito, Takeshima, Tadashi, Exploring suicide risk factors among Japanese individuals: the largest case-control psychological autopsy study in Japan. *Asian Journal of Psychiatry* <http://dx.doi.org/10.1016/j.ajp.2017.02.029>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Mental disorders and sleep problems are suicide risk factors.
- Verbal communication of one's own death is also a risk factor for suicide.
- They are suicide risk factors, independent of other factors.
- Forty-four percent of suicide completers saw a psychiatrist/psychosomatic doctor.

Download English Version:

<https://daneshyari.com/en/article/4930011>

Download Persian Version:

<https://daneshyari.com/article/4930011>

[Daneshyari.com](https://daneshyari.com)