

## Subject Index Volumes 72-79 (2017)

- Abandonment, related to history of suicide attempts among patients with major depressive disorders, early maladaptive schemas of, **77**, 71-79
- Acute inpatient care, dissociation in patients with borderline personality disorder in, **78**, 67-75
- AdAS Spectrum, *see* Adult Autism Subthreshold Spectrum
- ADHD, *see* Attention-deficit/hyperactivity disorder
- Adolescents
- and children, at clinical high risk for psychosis, suicidal behaviors and relationship with psychotic-like symptoms in, **78**, 31-37
  - and children, experiencing two major earthquakes in remote mountainous regions, mental health problems among, **72**, 66-73
  - and children, internalizing and externalizing disorders in, **75**, 75-84
  - mediational role of theory of mind in relation between attachment and thought problems in inpatient, **78**, 38-47
  - trauma type as conditional risk factor for posttraumatic stress disorder in referred clinic sample of, **76**, 138-146
  - and young adult, relatives of probands with bipolar disorder, substance use disorders in, **78**, 130-139
- Adult Autism Subthreshold Spectrum (AdAS Spectrum), questionnaire to assess subthreshold autism spectrum disorder, **73**, 61-83
- Adulthood, trait anxiety, through self-esteem, childhood parental bonding affects, **74**, 15-20
- Affect, positive and negative, to upper extremity physical function, emotion regulation strategies mediate associations of, **75**, 85-93
- Affective Neuroscience Personality Scale (ANPS), primary emotional traits and depressive tendencies, individual differences in, **73**, 136-142
- Age
- 1½–90+ years, empirically based assessment and taxonomy of psychopathology for, **79**, 4-18
  - early, of menarche, transdiagnostic psychiatric disorder risk associated with, **79**, 70-79
  - of onset, comorbidity, and suicidality, in obsessive-compulsive disorder, **76**, 79–86
  - and sex, gambling disorder associated with impulsivity traits measured by UPPS-P and association of, **72**, 106-113
- Aggression
- overt, and subjective anger, in psychiatric outpatients, **73**, 23-30
  - physical, and impulsive behaviors, in relation with BPD symptoms and criminal justice system involvement, role of emotion-driven difficulties controlling, **76**, 26-35
- Aggressive antisocial behavior, child, protective effect of character maturity in, **76**, 129-137
- Alcohol consumption, acute, impact on lethality of suicide methods, **75**, 27-34
- Alcohol dependence, effects on suicidality outcomes via both depression and posttraumatic stress in firefighters, **74**, 44-52
- Alcohol use, in low-risk population, pregnancy-specific anxiety associated with, **75**, 6-13
- Alexithymia
- and gastrointestinal-specific anxiety, role as predictors of treatment outcome in irritable bowel syndrome, **73**, 127-135
  - and non-suicidal self-injury, in borderline personality disorder, relationship between, **77**, 80-88
- Alzheimer's disease, and death, risk factors of transition from mild cognitive impairment to, **78**, 91-97
- Amyloid pathology, gender differences of neuropsychological profiles in cognitively normal older people without, **75**, 22-26
- Anger, subjective, and overt aggression, in psychiatric outpatients, **73**, 23-30
- Anhedonia, depression, and suicidal thoughts and behaviors, in adults with history of self-injury, relations between, **73**, 187-195
- ANPS, *see* Affective Neuroscience Personality Scale
- Antisocial personality disorder, neurocognitive dysfunction in problem gamblers with co-occurring, **76**, 153-159
- Anxiety
- bias in beliefs about self is associated with depression but not, **74**, 180-188
  - co-occurring, and trichotillomania, **72**, 1-5
  - and depression, in nonpsychiatric clinical settings, HEI for screening of, **76**, 87-97
  - depression and stress, impulsivity and its relationship with, **74**, 173-179
  - pregnancy-specific, associated with smoking, alcohol use and weight gain in low-risk population, **75**, 6-13
  - sensitivity, role in reactivity to trauma cues in treatment-seeking adults with substance use disorders, **78**, 107-114
- Anxious distress specifier interview, DSM-5, reliability and validity of, **76**, 11-17
- Attachment
- insecurity, self-reported, maternal oxytocin, and depressive symptoms with maternal caregiving sensitivity, associations of, **76**, 56-68
  - and thought problems, in inpatient adolescents, mediational role of theory of mind in relation between, **78**, 38-47
- Attention, to emotion through go/no-go task in children with oppositionality and callous–unemotional traits, **75**, 35-45
- Attention-deficit/hyperactivity disorder
- adult in comparison with borderline personality disorder and control participants, insomnia in, **76**, 119-128
  - and conduct disorder, neurocognitive characteristics of youth with noncomorbid and comorbid forms of, **77**, 60-70
  - dynamic thiol/disulfide homeostasis in children with, **73**, 53-60
  - phenomenology of hoarding in children with comorbid, **76**, 1-10

- Attentional bias  
 assessed by online shopping-related Stroop in high online shopping addiction tendency college students, self-control predicts, **75**, 14-21  
 in BDD via emotional Stroop task with eye-tracking paradigm, **74**, 151-161
- Attentional impulsivity, interactively predict food addiction in obese individuals, **72**, 83-87
- Attitude, towards suicide, preliminary evaluation of, **72**, 56-65
- Auditory verbal hallucinations, role of set-shifting in, **74**, 162-172
- Autism spectrum disorder, AdAS Spectrum questionnaire to assess subthreshold, **73**, 61-83
- Autoimmune encephalitis, and psychosis, clinical characterization of, **74**, 9-14
- BDD, *see* Body dysmorphic disorder
- BDNF, *see* Brain-derived neurotrophic factor
- Behavioral problems, influence on repetitive nonsuicidal self-injury in school sample, **74**, 214-223
- Beliefs, about self is associated with depression but not anxiety, bias in, **74**, 180-188
- Bifactor model, of comorbidity, temporal stability of, **72**, 74-82
- Binge eating disorder, comprehensive model of food addiction in patients with, **74**, 118-124
- Biological parents, insight into their offspring's schizophrenia associated with cognitive functioning and expressed emotion, **76**, 98-103
- Biomolecular aspects, of depression, **73**, 168-180
- Bipolar disorder  
 associated with less favorable socio-demographic and certain clinical features, lifetime presence of psychotic symptoms in, **76**, 169-176  
 exploration of metacognitive beliefs and thought control strategies in, **73**, 84-92  
 in manic phase, decreased S100B serum levels after treatment in patients with, **74**, 27-34  
 purinergic system dysfunctions in, **73**, 1-6  
 substance use disorders in adolescent and young adult relatives of probands with, **78**, 130-139
- Bipolar I disorder  
 with comorbid PTSD, **72**, 13-17  
 in remission, *vs.* schizophrenia in remission, **72**, 130-135
- Bipolar spectrum, two-dimensional model, integrating DSM-5's mixed features specifier for major depressive disorder, validation of, **77**, 89-99
- Body dysmorphic disorder  
 attentional biases in, via emotional Stroop task with eye-tracking paradigm, **74**, 151-161  
 relative to obsessive-compulsive disorder and psychotic disorders for DSM-5, insight in, **77**, 100-108
- Body image, in patients with mental disorders, **74**, 53-60
- Borderline personality disorder  
 in acute inpatient care, dissociation in patients with, **78**, 67-75  
 and control participants, insomnia in adult attention-deficit/hyperactivity disorder in comparison with, **76**, 119-128  
 relationship between non-suicidal self-injury and alexithymia in, **77**, 80-88  
 role of emotion-driven difficulties controlling impulsive behaviors and physical aggression in relation with symptoms of, **76**, 26-35
- Brain-derived neurotrophic factor  
 serum, and depression severity, improved by using HAM-D6, relationship between, **75**, 46-52
- Callous-unemotional traits, attention to emotion through go/no-go task in children with oppositionality and, **75**, 35-45
- Cardiometabolic risk, and sedentary behavior, among inpatients with schizophrenia, physical activity ameliorates association between, **74**, 144-150
- Cardiovascular risk, and metabolic syndrome, among institutionalized patients with schizophrenia receiving long term tertiary care, **74**, 196-203
- Caregiver, emotional intelligence and coping strategies as determinants of quality of life in depressed patient, **74**, 70-79
- Character maturity, protective effect in child aggressive antisocial behavior, **76**, 129-137
- Childhood adversity, self-reported, in relationship between patients with schizophrenia, **72**, 48-55
- Childhood trauma  
 effect on cognitive functions in sample of Chinese patients with schizophrenia, **76**, 147-152  
 subtypes, on fatigue and physical functioning in chronic fatigue syndrome, differential effects of, **78**, 76-82
- Children  
 and adolescents, at clinical high risk for psychosis, suicidal behaviors and relationship with psychotic-like symptoms in, **78**, 31-37  
 and adolescents, experiencing two major earthquakes in remote mountainous regions, mental health problems among, **72**, 66-73  
 and adolescents, internalizing and externalizing disorders in, **75**, 75-84  
 aggressive antisocial behavior, protective effect of character maturity in, **76**, 129-137  
 with attention deficit hyperactivity disorder, dynamic thiol/disulfide homeostasis in, **73**, 53-60  
 with comorbid ADHD, phenomenology of hoarding in, **76**, 1-10  
 in Lithuania, risk factors for mental health problems in, **73**, 15-22  
 with oppositionality and callous-unemotional traits, attention to emotion through go/no-go task in, **75**, 35-45  
 parental bonding, affects adulthood trait anxiety through self-esteem, **74**, 15-20  
 relationships among recollections of maternal and paternal acceptance–rejection in, and level of psychological adjustment among adults diagnosed with schizophrenia, social anxiety, and nonclinical control, **77**, 27-37
- Chronic diseases, suicidal ideation and suicidal attempts among adults with, **73**, 160-167
- Chronic fatigue syndrome, differential effects of childhood trauma subtypes on fatigue and physical functioning in, **78**, 76-82
- Clinical disorders, in outpatient psychiatric sample, joint hierarchical structure of, **79**, 19-30
- Clinicians  
 conflicting emotional responses to high suicide-risk patients and short-term suicide behaviors, **76**, 69-78  
 on Yale–Brown Obsessive Compulsive Scale, symptom correspondence between patients and, **73**, 105-110
- Cognitive-behavioral therapy, -based treatment were investigated in outpatients with eating disorders, psychological well-being changes following, **74**, 61-69
- Cognitive failures, in high schizotypy relate to emotional working memory deficits in lab, **78**, 115-129
- Cognitive functions  
 and expressed emotion, biological parents insight into their offspring's schizophrenia associated with, **76**, 98-103  
 in sample of Chinese patients with schizophrenia, effect of childhood trauma on, **76**, 147-152
- Cognitive impairments, among patients with major depressive disorder, elevated triglyceride levels associated with, **75**, 103-109
- Cognitive performance, in patients with schizophrenia, serum levels of interleukin-33 and soluble form receptor (sST2) associated with, **74**, 96-101
- College students, Chinese, exploring depression, self-esteem and verbal fluency with different degrees of internet addiction among, **72**, 114-120

Download English Version:

<https://daneshyari.com/en/article/4930168>

Download Persian Version:

<https://daneshyari.com/article/4930168>

[Daneshyari.com](https://daneshyari.com)