## Accepted Manuscript

Two distinct classes of degenerative change are independently linked to clinical progression in Mild Cognitive Impairment

Jean-Philippe Coutu, PhD, Emily R. Lindemer, PhD, Ender Konukoglu, PhD, David H. Salat, PhD

PII: S0197-4580(17)30040-4

DOI: 10.1016/j.neurobiolaging.2017.02.005

Reference: NBA 9843

To appear in: Neurobiology of Aging


Received Date: 2 May 2016

Revised Date: 15 January 2017

Accepted Date: 6 February 2017

Please cite this article as: Coutu, J.-P., Lindemer, E.R., Konukoglu, E., Salat, D.H., and the Alzheimer's Disease Neuroimaging Initiative (ADNI), Two distinct classes of degenerative change are independently linked to clinical progression in Mild Cognitive Impairment, *Neurobiology of Aging* (2017), doi: 10.1016/j.neurobiolaging.2017.02.005.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## Two distinct classes of degenerative change are independently linked to clinical progression in Mild Cognitive Impairment

**Authors:** Jean-Philippe Coutu, PhD<sup>a,b,c,1</sup>\*; Emily R. Lindemer, PhD<sup>a,b,c</sup>, Ender Konukoglu, PhD<sup>a,c,2</sup>; David H. Salat, PhD<sup>a,c,d</sup> and the Alzheimer's Disease Neuroimaging Initiative (ADNI)†

a. MGH/MIT/HMS Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Harvard Medical School, Charlestown, MA, USA; b. Harvard-MIT Division of Health Sciences and Technology, Massachusetts Institute of Technology, Cambridge, MA, USA; c. Departments of Radiology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA; d. Neuroimaging Research for Veterans Center, VA Boston Healthcare System, Boston, MA, USA.

<sup>†</sup>Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at <u>http://adni.loni.usc.edu/wp-</u>

content/uploads/how\_to\_apply/ADNI\_Acknowledgement\_List.pdf

**Corresponding author:** Jean-Philippe Coutu; <u>coutu@nmr.mgh.harvard.edu</u>; Address: Athinoula A. Martinos Center for Biomedical Imaging, 149 13<sup>th</sup> Street, Room 2301, Charlestown, MA 02129, USA; Phone: 1-857-318-9617; Fax: 1-617-726-7422.

<sup>&</sup>lt;sup>1</sup> Biospective Inc, Montreal, Canada

<sup>&</sup>lt;sup>2</sup> Computer Vision Laboratory, ETH Zurich, Switzerland

Download English Version:

## https://daneshyari.com/en/article/4932543

Download Persian Version:

https://daneshyari.com/article/4932543

Daneshyari.com