

Author's Accepted Manuscript

Associations between self-harm and distinct types of impulsivity

Samuel R. Chamberlain, Sarah A. Redden, Jon E. Grant



PII: S0165-1781(16)31021-6
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.01.050>
Reference: PSY10247

To appear in: *Psychiatry Research*

Received date: 13 June 2016
Revised date: 3 December 2016
Accepted date: 20 January 2017

Cite this article as: Samuel R. Chamberlain, Sarah A. Redden and Jon E. Grant Associations between self-harm and distinct types of impulsivity, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.01.050>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and a review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Associations between self-harm and distinct types of impulsivity^{☆☆}

Samuel R Chamberlain MD PhD^{a,b}, Sarah A. Redden BA^c, Jon E. Grant JD MD MPH^{c*}

^aDepartment of Psychiatry, University of Cambridge, UK

^bCambridge and Peterborough NHS Foundation Trust, UK

^cDepartment of Psychiatry & Behavioral Neuroscience, University of Chicago

* **Correspondence to:** Jon E. Grant, JD, MD, MPH Professor, Department of Psychiatry & Behavioral Neuroscience University of Chicago, Pritzker School of Medicine 5841 S. Maryland Avenue, MC 3077, Chicago, IL 60637. Phone: 773-834-1325; Fax: 773-834-6761.

jongrant@uchicago.edu

ABSTRACT**Objective**

There is an ongoing debate regarding how self-harm should be classified. The aim of this study was to characterize associations between self-harm and impulsivity.

Method

Total 333 adults (mean [SD] age 22.6 (3.6) years, 61% male) were recruited from the general community. History of self-harm was quantified using the Self-Harm Inventory (SHI), which asks about 22 self-harm behaviors. Principal components analysis was used to identify latent dimensions of self-harming behaviors. Relationships between self-harm dimensions and other measures were characterized using ordinary least squares regression.

[☆] Authors' version of resubmitted manuscript, Psych Res 2016

[☆] **Previous presentations:** none.

Download English Version:

<https://daneshyari.com/en/article/4933330>

Download Persian Version:

<https://daneshyari.com/article/4933330>

[Daneshyari.com](https://daneshyari.com)