


REVERSAL LEARNING IN PATIENTS WITH
OBSESSIVE-COMPULSIVE DISORDER (OCD)
AND THEIR UNAFFECTED RELATIVES: IS
ORBITOFRONTAL DYSFUNCTION AN
ENDOPHENOTYPE OF OCD?

Didem Tezcan, Selim Tumkaya, Emre Bora



PII: S0165-1781(16)31267-7
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.03.001>
Reference: PSY10368

To appear in: *Psychiatry Research*

Received date: 1 August 2016
Revised date: 26 November 2016
Accepted date: 1 March 2017

Cite this article as: Didem Tezcan, Selim Tumkaya and Emre Bora, REVERSAL LEARNING IN PATIENTS WITH OBSESSIVE-COMPULSIVE DISORDER (OCD) AND THEIR UNAFFECTED RELATIVES: IS ORBITOFRONTAL DYSFUNCTION AN ENDPHENOTYPE OF OCD?, *Psychiatry Research* <http://dx.doi.org/10.1016/j.psychres.2017.03.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

REVERSAL LEARNING IN PATIENTS WITH OBSESSIVE-
COMPULSIVE DISORDER (OCD) AND THEIR UNAFFECTED
RELATIVES: IS ORBITOFRONTAL DYSFUNCTION AN
ENDOPHENOTYPE OF OCD?

Didem Tezcan^a, Selim Tumkaya^{a*}, Emre Bora^b

^aPamukkale University, Medicine Faculty, Department of Psychiatry, Denizli,
(20070), Turkey.

^bMelbourne Neuropsychiatry Centre, Department of Psychiatry. University of
Melbourne, VIC, Australia; Department of Psychiatry, University of Dokuz
Eylül, İzmir, Turkey.

* Corresponding author. Selim Tumkaya Pamukkale University, Medicine
Faculty, Department of Psychiatry, Kınıklı, Denizli (20070), Turkey. Tel.: +90
4440728-5016; fax: +90(258)2134922; selimtumkaya@gmail.com

SUMMARY

It has been suggested that reversal learning deficits might be an endophenotype of OCD. To investigate this hypothesis, we administered a probabilistic reversal learning task (ProbRev) to OCD patients, their unaffected first-degree relatives, and healthy controls. Although the relatives had a performance in between OCDs and controls at the early

Download English Version:

<https://daneshyari.com/en/article/4933443>

Download Persian Version:

<https://daneshyari.com/article/4933443>

[Daneshyari.com](https://daneshyari.com)