

Anticholinergic use trends in 14,013 patients with schizophrenia from three national surveys on the use of psychotropic medications in China (2002-2012)

Yun-Ai Su, Feng Yan, Qian Li, Yu-Tao Xiang, Liang Shu, Xin Yu, Yu-Ping Ning, Ke-Rang Zhang, Tao Li, Qi-Yi Mei, Ke-Qing Li, Tian-Mei Si



PII: S0165-1781(16)32103-5
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.07.038>
Reference: PSY10672

To appear in: *Psychiatry Research*

Received date: 12 December 2016
Revised date: 25 April 2017
Accepted date: 21 July 2017

Cite this article as: Yun-Ai Su, Feng Yan, Qian Li, Yu-Tao Xiang, Liang Shu, Xin Yu, Yu-Ping Ning, Ke-Rang Zhang, Tao Li, Qi-Yi Mei, Ke-Qing Li and Tian-Mei Si, Anticholinergic use trends in 14,013 patients with schizophrenia from three national surveys on the use of psychotropic medications in China (2002-2012), *Psychiatry Research* <http://dx.doi.org/10.1016/j.psychres.2017.07.038>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Anticholinergic use trends in 14,013 patients with schizophrenia from three national surveys on the use of psychotropic medications in China (2002-2012)

Yun-Ai Su^{a, 1}, Feng Yan^{a, b, 1}, Qian Li^a, Yu-Tao Xiang^{c, d}, Liang Shu^a, Xin Yu^a, Yu-Ping Ning^e, Ke-Rang Zhang^f, Tao Li^g, Qi-Yi Mei^h, Ke-Qing Liⁱ, Tian-Mei Si^{a*}

- a. Peking University Sixth Hospital (Institute of Mental Health) & National Clinical Research Center for Mental Disorders & Key Laboratory of Mental Health, Ministry of Health (Peking University), Beijing, China;
- b. Beijing Hui-Long-Guan Hospital, Peking University, Beijing, China;
- c. The National Clinical Research Center for Mental Disorders, China & Center of Depression, Beijing Institute for Brain Disorders & Beijing Anding Hospital, Capital Medical University, Beijing, China;
- d. Unit of Psychiatry, Faculty of Health Sciences, University of Macau, Macao SAR, China;
- e. Psychiatric Hospital, Guangzhou, China;
- f. The First Hospital of Shanxi Medical University, Shanxi province, China;
- g. West China Hospital, Sichuan University, China;
- h. Suzhou Guangji Hospital, Suzhou, China;
- i. Hebei Mental Health Center, Hebei, China

Corresponding Authors:

Tian-Mei Si, National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital/ Institute of Mental Health), and the Key Laboratory of Mental Health, Ministry of Health (Peking University), Beijing 100191, China; Tel.: +86 10 62723748; Fax: +86 10 62027314; E-mail address: si.tian-mei@163.com (T.-M. Si);

¹These authors contributed equally to this work.

Download English Version:

<https://daneshyari.com/en/article/4933470>

Download Persian Version:

<https://daneshyari.com/article/4933470>

[Daneshyari.com](https://daneshyari.com)