

Author's Accepted Manuscript

Acute effects of smoked marijuana in marijuana smokers at clinical high-risk for psychosis: a preliminary study

Nehal P. Vadhan, Cheryl M. Corcoran, Gill Bedi, John G Keilp, Margaret Haney



PII: S0165-1781(17)30158-0
DOI: <http://dx.doi.org/10.1016/j.psychres.2017.07.070>
Reference: PSY10704

To appear in: *Psychiatry Research*

Received date: 27 January 2017
Revised date: 8 June 2017
Accepted date: 30 July 2017

Cite this article as: Nehal P. Vadhan, Cheryl M. Corcoran, Gill Bedi, John G Keilp and Margaret Haney, Acute effects of smoked marijuana in marijuana smokers at clinical high-risk for psychosis: a preliminary study, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.07.070>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Acute effects of smoked marijuana in marijuana smokers at clinical high-risk for psychosis: a preliminary study

Nehal P. Vadhan^{a*}, Cheryl M. Corcoran^b, Gill Bedi^b, John G Keilp^b & Margaret Haney^b

^aDepartments of Psychiatry and Molecular Medicine, Hofstra Northwell School of Medicine and the Center for Psychiatric Neuroscience, Feinstein Institute for Medical Research, Great Neck, NY, USA

^bDepartment of Psychiatry, Columbia University Medical Center and the New York State Psychiatric Institute, New York, NY, USA

1.1 *Corresponding author. Nehal P. Vadhan, PhD, Department of Psychiatry, Northwell Health, 1010 Northern Blvd, Suite #311, Great Neck, NY 11021, P: 516-837-1685; F: 516-837-1699, E: nvadhan@northwell.edu

Abstract

Marijuana use is associated with psychosis, but its effects are understudied in individuals with preexisting risk for psychotic disorders. This preliminary study examined the acute psychological and physiological effects of smoked marijuana (0.0% or 5.5% Δ^9 -THC) in marijuana users at clinical high-risk (CHR; n=6) to develop a psychotic disorder, and those not at risk (n=6), under controlled laboratory

Download English Version:

<https://daneshyari.com/en/article/4933535>

Download Persian Version:

<https://daneshyari.com/article/4933535>

[Daneshyari.com](https://daneshyari.com)