

Author's Accepted Manuscript

Thwarted Interpersonal Needs and Suicide Ideation: Comparing Psychiatric Inpatients with Bipolar and Non-Bipolar Mood Disorders

Nathanael J. Taylor, Sean M. Mitchell, Jared F. Roush, Sarah L. Brown, Danielle R. Jahn, Kelly C. Cukrowicz



PII: S0165-1781(16)30217-7
DOI: <http://dx.doi.org/10.1016/j.psychres.2016.09.025>
Reference: PSY9959

To appear in: *Psychiatry Research*

Received date: 5 February 2016
Revised date: 29 August 2016
Accepted date: 18 September 2016

Cite this article as: Nathanael J. Taylor, Sean M. Mitchell, Jared F. Roush, Sarah L. Brown, Danielle R. Jahn and Kelly C. Cukrowicz, Thwarted Interpersonal Needs and Suicide Ideation: Comparing Psychiatric Inpatients with Bipolar and Non-Bipolar Mood Disorders, *Psychiatry Research* <http://dx.doi.org/10.1016/j.psychres.2016.09.025>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

Thwarted Interpersonal Needs and Suicide Ideation: Comparing Psychiatric Inpatients with
Bipolar and Non-Bipolar Mood Disorders

Nathanael J. Taylor^a, Sean M. Mitchell^a, Jared F. Roush^a, Sarah L. Brown^a, Danielle R. Jahn^b,

Kelly C. Cukrowicz^{a*}

^aDepartment of Psychological Sciences, Mail Stop 2051, Texas Tech University, Lubbock, TX
79409-2051

^bThe VA Capitol Health Care Network (VISN 5) Mental Illness Research, Education, and
Clinical Center; Department of Psychiatry, University of Maryland School of Medicine

*Corresponding author. Department of Psychological Sciences, Mail Stop 42051, Texas Tech
University, Lubbock, TX 79409-2051. Tel.: 806 834 8485, fax: 806 742 0818.

kelly.cukrowicz@ttu.edu

Abstract

Psychiatric inpatients are at heightened risk for suicide, and evidence suggests that psychiatric inpatients with bipolar mood disorders may be at greater risk for suicide ideation compared to those with non-bipolar mood disorders. There is a paucity of research directly comparing risk factors for suicide ideation in bipolar versus non-bipolar mood disorders in an inpatient sample. The current study sought to clarify the role of two constructs from the interpersonal theory of suicide (i.e., perceived burdensomeness and thwarted belongingness) in leading to suicide ideation among psychiatric inpatients with bipolar and non-bipolar mood disorders. Participants were ($N = 90$) psychiatric inpatients with a bipolar ($n = 20$) or non-bipolar mood disorder ($n = 70$; per their medical charts). Perceived burdensomeness, but not thwarted

Download English Version:

<https://daneshyari.com/en/article/4933777>

Download Persian Version:

<https://daneshyari.com/article/4933777>

[Daneshyari.com](https://daneshyari.com)