

Author's Accepted Manuscript

Maladaptive Five Factor Model Personality Traits Associated with Borderline Personality Disorder Indirectly Affect Susceptibility to Suicide Ideation Through Increased Anxiety Sensitivity Cognitive Concerns

Raymond P. Tucker, Greg J. Lengel, Caitlin E. Smith, Dan W. Capron, Stephanie N. Mullins-Sweatt, LaRicka R. Wingate



PII: S0165-1781(15)30722-8
DOI: <http://dx.doi.org/10.1016/j.psychres.2016.08.051>
Reference: PSY9922

To appear in: *Psychiatry Research*

Received date: 22 November 2015
Accepted date: 17 August 2016

Cite this article as: Raymond P. Tucker, Greg J. Lengel, Caitlin E. Smith, Dan W. Capron, Stephanie N. Mullins-Sweatt and LaRicka R. Wingate, Maladaptive Five Factor Model Personality Traits Associated with Borderline Personality Disorder Indirectly Affect Susceptibility to Suicide Ideation Through Increased Anxiety Sensitivity Cognitive Concerns, *Psychiatry Research* <http://dx.doi.org/10.1016/j.psychres.2016.08.051>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

Maladaptive Five Factor Model Personality Traits Associated with Borderline Personality
Disorder Indirectly Affect Susceptibility to Suicide Ideation Through Increased Anxiety
Sensitivity Cognitive Concerns

Raymond P. Tucker^a, Greg J. Lengel^a, Caitlin E. Smith^a, Dan W. Capron^b, Stephanie N. Mullins-
Sweatt^a, LaRicka R. Wingate^{a*}

^aOklahoma State University: 116 North Murray, USA, Stillwater, OK 74078

^bUniversity of Southern Mississippi 118 College Drive #5025, USA, Hattiesburg, MS 39406-
5025

*Correspondence may be sent to: Oklahoma State University, Department of Psychology, 116 North Murray Hall, USA, Stillwater, OK 74078. Tel.: +405-744-5543. fax: 405-744-8067. laricka.wingate@okstate.edu.

1. Introduction

BPD is a significant clinical syndrome characterized by heightened emotion dysregulation, impulsivity, rocky and unstable interpersonal relationships, and chronic feelings of emptiness. Suicidal behavior and suicide ideation are common experiences for those with BPD and are included as symptoms of the disorder in the Diagnostic and Statistical Manual of Mental Disorders-5 (DSM-5; APA, 2013). Psychological autopsy studies have suggested that up to 33% of participants who died by suicide might have been diagnosed with BPD if they had been assessed (Kullgren et al., 1986; Runeson and Beskow, 1991). The risk for attempting suicide is significantly increased in psychiatric inpatients diagnosed with an affective disorder (e.g., major depressive disorder, schizoaffective disorder, and bipolar disorder) when BPD is

Download English Version:

<https://daneshyari.com/en/article/4933815>

Download Persian Version:

<https://daneshyari.com/article/4933815>

[Daneshyari.com](https://daneshyari.com)