


# Author's Accepted Manuscript

Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing

Emma Komulainen, Enrico Glerean, Katarina Meskanen, Roope Heikkilä, Lauri Nummenmaa, Tuukka T. Raij, Jari Lahti, Pekka Jylhä, Tarja Melartin, Erkki Isometsä, Jesper Ekelund



PII: S0925-4927(16)30336-5  
DOI: <http://dx.doi.org/10.1016/j.psychresns.2017.03.009>  
Reference: PSYN10663

To appear in: *Psychiatry Research: Neuroimaging*

Received date: 13 November 2016

Revised date: 17 February 2017

Accepted date: 20 March 2017

Cite this article as: Emma Komulainen, Enrico Glerean, Katarina Meskanen, Roope Heikkilä, Lauri Nummenmaa, Tuukka T. Raij, Jari Lahti, Pekka Jylhä, Tarja Melartin, Erkki Isometsä and Jesper Ekelund, Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing, *Psychiatry Research: Neuroimaging* <http://dx.doi.org/10.1016/j.psychresns.2017.03.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and a review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing**

Emma Komulainen<sup>a</sup>, Enrico Glerean<sup>b</sup>, Katarina Meskanen<sup>a</sup>, Roope Heikkilä<sup>a</sup>, Lauri Nummenmaa<sup>b,c</sup>, Tuukka T. Raij<sup>a,b,d</sup>, Jari Lahti<sup>e,f,g</sup>, Pekka Jylhä<sup>a,h</sup>, Tarja Melartin<sup>a</sup>, Erkki Isometsä<sup>a,h</sup>, Jesper Ekelund<sup>a,i</sup>.

<sup>a</sup>University of Helsinki and Helsinki University Hospital, Psychiatry, Helsinki, Finland

<sup>b</sup>Aalto University, School of Science, Department of Neuroscience and Biomedical Engineering, Espoo, Finland

<sup>c</sup>Turku PET Centre and Department of Psychology, University of Turku, Finland

<sup>d</sup>Aalto NeuroImaging, Aalto University, Espoo, Finland

<sup>e</sup>University of Helsinki, Institute of Behavioral Sciences, Helsinki, Finland

<sup>f</sup>Folkhälsan Research Center, Helsinki, Finland

<sup>g</sup>Helsinki collegium of Advanced Studies, University of Helsinki, Finland

<sup>h</sup>National Institute of Health and Welfare, Department of Mental Health and Substance Abuse Services, Helsinki, Finland

<sup>i</sup>Vaasa Hospital District, Department of Psychiatry, Vaasa, Finland

Corresponding author:

Emma Komulainen

Department of Psychiatry, University of Helsinki, P.O. Box 590, FI-00029 HUS, Finland

E-mail: emma.komulainen@helsinki.fi

Phone: +358505203573

Fax number: +358947163735

**Abstract**

The link between neurotransmitter-level effects of antidepressants and their clinical effect remain poorly understood. A single dose of mirtazapine decreases limbic responses to fearful faces in

Download English Version:

<https://daneshyari.com/en/article/4933944>

Download Persian Version:

<https://daneshyari.com/article/4933944>

[Daneshyari.com](https://daneshyari.com)