

Accepted Manuscript

Title: Effects of Antiepileptic Drugs on Thyroid Hormone Function in Epilepsy Patients

Authors: Fu-Yuan Shih, Yao-Chung Chuang, Ming-Jung Chuang, Yan-Ting Lu, Wan-Chen Tsai, Ting-Ying Fu, Meng-Han Tsai



PII: S1059-1311(17)30194-2
DOI: <http://dx.doi.org/doi:10.1016/j.seizure.2017.03.011>
Reference: YSEIZ 2918

To appear in: *Seizure*

Received date: 17-10-2016
Revised date: 17-3-2017
Accepted date: 18-3-2017

Please cite this article as: Shih Fu-Yuan, Chuang Yao-Chung, Chuang Ming-Jung, Lu Yan-Ting, Tsai Wan-Chen, Fu Ting-Ying, Tsai Meng-Han. Effects of Antiepileptic Drugs on Thyroid Hormone Function in Epilepsy Patients. *SEIZURE: European Journal of Epilepsy* <http://dx.doi.org/10.1016/j.seizure.2017.03.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Effects of Antiepileptic Drugs on Thyroid Hormone Function in Epilepsy Patients

Fu-Yuan Shih^{1†}, Yao-Chung Chuang^{2,3,4,5†}, Ming-Jung Chuang¹, Yan-Ting Lu², Wan-Chen Tsai²,

Ting-Ying Fu⁶, and Meng-Han Tsai^{2,7*} (✉)

Correspondence to:

Meng-Han Tsai, MD, PhD

Department of Neurology, Kaohsiung Chang Gung Memorial Hospital,

123, Dapi Road, Niasong District, Kaohsiung 83304, Taiwan

Tel: +886-7-7317123 ext. 2285

E-mail: menghan@cgmh.org.tw

†Contributed equally to the manuscript

¹ Departments of Neurosurgery, Kaohsiung Chang Gung Memorial Hospital, Chang Gung University College of Medicine, Kaohsiung, Taiwan.

² Department of Neurology, Kaohsiung Chang Gung Memorial Hospital, Chang Gung University College of Medicine, Kaohsiung, Taiwan.

³ Center for Translational Research in Biomedical Sciences, Kaohsiung, Taiwan.

⁴ Department of Biological Science, National Sun Yet-Sen University, Kaohsiung, Taiwan.

⁵ Faculty of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

⁶ Department of Pathology and Laboratory Medicine, Kaohsiung Veteran General Hospital, Kaohsiung, Taiwan.

⁷ Department of Computer Science and Information Engineering, Cheng Shiu University, Kaohsiung, Taiwan

Highlights

- 17.4% epilepsy patients had low fT4.
- Female patients with epilepsy and an older age, AED polytherapy, and carbamazepine treatment had a higher risk of low fT4.
- Thyroid function in elderly female epilepsy patients who received carbamazepine therapy should be monitored.

Download English Version:

<https://daneshyari.com/en/article/4935411>

Download Persian Version:

<https://daneshyari.com/article/4935411>

[Daneshyari.com](https://daneshyari.com)