

# Accepted Manuscript

High order spectral difference lattice Boltzmann method for incompressible hydrodynamics

Weidong Li

PII: S0021-9991(17)30422-9  
DOI: <http://dx.doi.org/10.1016/j.jcp.2017.05.039>  
Reference: YJCPH 7386

To appear in: *Journal of Computational Physics*

Received date: 7 October 2016  
Revised date: 17 April 2017  
Accepted date: 22 May 2017

Please cite this article in press as: W. Li, High order spectral difference lattice Boltzmann method for incompressible hydrodynamics, *J. Comput. Phys.* (2017), <http://dx.doi.org/10.1016/j.jcp.2017.05.039>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## Highlights

- High order spectral difference lattice Boltzmann method (SD-LBM) is proposed.
- SD-LBM can be implemented on arbitrary unstructured quadrilateral meshes.
- SD-LBM almost holds all advantages of the LBM, but is more flexible.
- SD-LBM is simple and accurate due to only first order derivatives involved in LBE.

Download English Version:

<https://daneshyari.com/en/article/4967423>

Download Persian Version:

<https://daneshyari.com/article/4967423>

[Daneshyari.com](https://daneshyari.com)