## Accepted Manuscript

#### Regular Article

Metallogel Templeted Synthesis and Stabilization of Silver-Particles and Its Application in Catalytic Reduction of Nitro-Arene

Mukesh Sharma, Palaban Jyoti Sarma, Manash Goswami, Kusum K. Bania

PII:	S0021-9797(16)30942-0
DOI:	http://dx.doi.org/10.1016/j.jcis.2016.11.065
Reference:	YJCIS 21793
To appear in:	Journal of Colloid and Interface Science
Received Date:	23 September 2016
Revised Date:	16 November 2016
Accepted Date:	19 November 2016



Please cite this article as: M. Sharma, P. Jyoti Sarma, M. Goswami, K.K. Bania, Metallogel Templeted Synthesis and Stabilization of Silver-Particles and Its Application in Catalytic Reduction of Nitro-Arene, *Journal of Colloid and Interface Science* (2016), doi: http://dx.doi.org/10.1016/j.jcis.2016.11.065

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# ACCEPTED MANUSCRIPT

## Metallogel Templeted Synthesis and Stabilization of Silver-Particles and Its Application in Catalytic Reduction of Nitro-Arene

Mukesh Sharma,<sup>a</sup> Palaban Jyoti Sarma,<sup>a</sup> Manash Goswami<sup>b</sup> and Kusum K. Bania<sup>a</sup>\*

<sup>a</sup>Department of Chemical Sciences, Tezpur University, Assam, India 784028 <sup>b</sup>Department of Chemistry, Gauhati University, Gopinath Bordoloi Nagar, Jalukbari, Guwahati, Assam 781014

\* Corresponding author at Department of Chemical Sciences, Tezpur University, Assam, India 784028, E-mail: <u>kusum@tezu.ernet.in</u>

### ABSTRACT

Metallogel of iron-carboxylates was obtained from trans 1,2 cyclohexanedicarboxylic acid in dimethylformamide (DMF) at basic condition. Spectroscopic and SEM morphology study of the iron-metallogel revealed that the iron complex with dicarboxylic acid was linked together via carboxylates and led to a supramolecular helical like architecture. The synthesized metallogel served as an excellent template for *in-situ* reduction of silver ion to silver particles micro to nano scale range. Variation of  $AgNO_3$  concentration shepherd to change the morphology of the Ag-particles. AgNO<sub>3</sub> concentration was found to affect the shape and size of silver particles. On going from lower to higher concentration shape of silver particles changed from spherical to large agglomerated particles. Cubic shape Ag-particles were found on treatment of 0.05M AgNO<sub>3</sub> solution with metallogel. Cubical shape silver particles were found to be effective catalyst for nitro-arene reduction in presence of NaBH<sub>4</sub>. Density functional theory (DFT) calculations were performed to rationalize the role of Ag-particles in catalytic reduction of 4-nitrophenol to 4-aminophenol. Based on DFT study, we proposed that catalytic reduction occurred via Ag-hydride complex formation. Since metallogels as well as the 4-aminophenol are finding large application in pharamaceuticals industries therefore the current work can provide an alternatives path in production of 4-aminophenols. In addition to this, the synthesis of Ag-nanomaterials using metallogel as template can pave a new direction in the development of nanotechnology and might find wide applications in catalytic industrial processes.

**Keywords**: *trans 1,2* cyclohexanedicarboxylic, DFT, Fe-Metallogel, Reduction of 4nitrophenol, Silver particles Download English Version:

https://daneshyari.com/en/article/4985214

Download Persian Version:

https://daneshyari.com/article/4985214

Daneshyari.com