Accepted Manuscript


Temporal pattern in Corinth rift seismicity revealed by visibility graph analysis

George Hloupis

 PII:
 S1007-5704(17)30093-X

 DOI:
 10.1016/j.cnsns.2017.03.015

 Reference:
 CNSNS 4142



To appear in: Communications in Nonlinear Science and Numerical Simulation

Received date:	6 September 2016
Revised date:	15 March 2017
Accepted date:	16 March 2017

Please cite this article as: George Hloupis, Temporal pattern in Corinth rift seismicity revealed by visibility graph analysis, *Communications in Nonlinear Science and Numerical Simulation* (2017), doi: 10.1016/j.cnsns.2017.03.015

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Visibility Graph (VG) method was applied for the 2001-2011 seismicity of Corinth rift.
- Using a sliding window approach the exponent γ of the degree distribution of VG was studied.
- A time coincidence exists between the minima of exponent γ and the period when the most significant seismic swarm took place.

Download English Version:

https://daneshyari.com/en/article/5011429

Download Persian Version:

https://daneshyari.com/article/5011429

Daneshyari.com