


Accepted Manuscript

Application of a Lie group admitted by a homogeneous equation for group classification of a corresponding inhomogeneous equation

Feng-Shan Long, Adisak Karnbanjong,
Amornrat Suriyawichitseranee, Yurii N. Grigoriev,
Sergey V. Meleshko

PII: S1007-5704(17)30003-5
DOI: [10.1016/j.cnsns.2017.01.006](https://doi.org/10.1016/j.cnsns.2017.01.006)
Reference: CNSNS 4077



To appear in: *Communications in Nonlinear Science and Numerical Simulation*

Received date: 17 December 2015
Revised date: 4 November 2016
Accepted date: 2 January 2017

Please cite this article as: Feng-Shan Long, Adisak Karnbanjong, Amornrat Suriyawichitseranee, Yurii N. Grigoriev, Sergey V. Meleshko, Application of a Lie group admitted by a homogeneous equation for group classification of a corresponding inhomogeneous equation, *Communications in Nonlinear Science and Numerical Simulation* (2017), doi: [10.1016/j.cnsns.2017.01.006](https://doi.org/10.1016/j.cnsns.2017.01.006)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- New method for group classification is given.
- The method can be applied for equations with nonlocal terms.
- The method is illustrated by several examples.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/5011605>

Download Persian Version:

<https://daneshyari.com/article/5011605>

[Daneshyari.com](https://daneshyari.com)