



# Spatial patterns of *Pisidium chilense* (Mollusca Bivalvia) and *Hyalella patagonica* (Crustacea, Amphipoda) in an unpolluted stream in Navarino island (54° S, Cape Horn Biosphere Reserve)

Patricio De Los Ríos Escalante<sup>a,b,\*</sup>, Andrés Mansilla<sup>c,d</sup>

<sup>a</sup> Laboratorio de Ecología Aplicada y Biodiversidad, Escuela de Ciencias Ambientales, Facultad de Recursos Naturales, Universidad Católica de Temuco, Casilla 15-D, Chile

<sup>b</sup> Núcleo Ambientales UCTemuco, Chile

<sup>c</sup> Departamento Ciencias y Recursos Naturales, Universidad de Magallanes, Av. Bulnes 01855, Punta Arenas, Chile

<sup>d</sup> Instituto de Ecología Aplicada y Biodiversidad, Casilla 653, Santiago de Chile, Chile

Received 24 May 2016; accepted 6 July 2016

Available online 4 August 2016

## KEYWORDS

*Pisidium chilense*;  
*Hyalella patagonica*;  
Species association;  
Negative binomial  
distribution

**Abstract** The southern South American inland waters have many endemic species and some of them are considered as endangered for IUCN, that inhabits in unpolluted ecosystems, one of these ecosystems are the sub-Antarctic perennial forests located in the Cape Horn Biosphere Reserve at 54° S. The aim of the present study is to analyze the spatial patterns of *Pisidium chilense* Ituarte, 1999 (Mollusca Bivalvia) and *Hyalella patagonica* (Cunningham, 1871) (Crustacea, Amphipoda) in an unpolluted stream. Both species had aggregated spatial distribution, both have a negative binomial distribution pattern, and both are associated. The present results would agree with similar patterns in Patagonian rivers where both species coexist.

© 2016 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The Southern Patagonia has unpolluted ecosystems with low human intervention due mainly to their geographic isolation and difficult access. There are many endemic species, especially in inland water bodies (Jara et al., 2006). The Cape Horn Biosphere Reserve is located in the southern island of South America with unpolluted and pristine sub-Antarctic perennial forests with unpolluted lakes, rivers and streams, the fauna is markedly endemic (Moorman et al., 2006). The benthic fauna

\* Corresponding author at: Laboratorio de Ecología Aplicada y Biodiversidad, Escuela de Ciencias Ambientales, Facultad de Recursos Naturales, Universidad Católica de Temuco, Casilla 15-D, Chile.

E-mail address: [prios@uct.cl](mailto:prios@uct.cl) (P. De Los Ríos Escalante).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

spatial distribution can be explained in accordance to probabilistic models that can explain if the species have random, aggregated or uniform spatial distribution (Elliot, 1983; Zar, 1999). On this basis, the spatial pattern can adjust to Poisson, negative binomial or binomial distribution respectively (Elliot, 1983; Fernandes et al., 2003; De los Rios-Escalante, 2011). The aim of the present study is to analyze the spatial distribution of two species representative from Southern Patagonian inland waters, the bivalve *Pisidium chilense* Ituarte, 1999 and the crustacean *Hyaella patagonica* (Cunningham, 1871), *P. chilense* is considered as endangered for IUCN (IUCN, 2015). Both species are important in inland water communities in Southern Patagonia (Moorman et al., 2006).


## 2. Materials and methods

The studied site was a small stream close to Robalo river located close to Puerto Williams town in an protected unpolluted area, close to Cape Horn Biosphere Reserve, in Navarino island (54° 55' 47,3' S; 67° 33' 43,8" W, Fig. 1). Samples were

collected from the riparian zone of this shallow stream using a Ø L plastic cube with a 1 mm mesh, 20 replicas were collected random along a transect of 50 m from the study site, and the specimens were counted *in situ*. The site was visited and sampled on the 10th February 2010. It was used in a 2 × 2 contingency table analysis with the species *H. patagonica* and *P. chilense* as the first and second categorical variables. The Mean abundance (and variance) was calculated for both species with the aim of determining if their spatial dispersions are aggregated, random or uniform using Variable/Mean ratio to determine if both species are associated (Zar, 1999; Gotelli and Ellison, 2004). Software Xlstat 10.0 was used when the population dispersions were described by the negative binomial distribution (Zar, 1999; Fernandes et al., 2003).

## 3. Results and discussion

The results of the 2 × 2 contingency table revealed first that both species are not independent in the sampled site, this means that both species share the habitat together ( $\chi^2$



**Figure 1** Location Navarino island (in gray), Puerto Williams town and photograph of studied site.

Download English Version:

<https://daneshyari.com/en/article/5023039>

Download Persian Version:

<https://daneshyari.com/article/5023039>

[Daneshyari.com](https://daneshyari.com)