



Contents lists available at ScienceDirect

Quaternary International

journal homepage: www.elsevier.com/locate/quaint

Changes in the fauna of bats in the south of the Russian Far East since the late Pleistocene



Mikhail P. Tiunov

Institute of Biology and Soil Science FEB RAS, Prospekt 100-letia 159, 690022 Vladivostok, Russia

ARTICLE INFO

Article history:

Received 18 May 2016

Received in revised form

27 September 2016

Accepted 29 September 2016

Keywords:

Bats

Far East Russia

Holocene

Late Pleistocene

ABSTRACT

The aim of the study was to identify changes in the fauna of bats during the late Pleistocene and Holocene in Southeast Russia, to utilize the results together with environmental data to disseminate modern analogues and reconstruct fossil environment. Fossil and subfossil lower jawbones of bats found in Medvezhyi Klyk cave deposits were identified by comparing morphology and surface features to modern species. The ecology and distribution of modern analogues are extrapolated to reconstruct the fossil environment. Fifteen bat species belonging to 6 genera were identified. Of the bat bone residues detected, the greatest number and the richest species composition formed approximately 40 ka BP in the lower layers of the cave sediments. In the layers formed in the Last Glacial Maximum period, the remains of only two species were found. The species diversity of bat bone residues gradually increased, reaching a maximum (7 species) in the layers corresponding to the optimum of the Holocene. At present, only single individuals of *Murina hilgendorfi* and *Plecotus ognevi* can occasionally be found hibernating in caves. Fossil remains of forest species are present in all layers of sediments, but occasionally, the remains of species of open landscapes not currently living in the territory appear as well. Studies have shown that during the warmer periods of the late Pleistocene and Holocene the species composition of bats in southern Primorye was expanded by southern thermophilic species that moved much farther north than previously thought. The presence of fauna of that time, including forest species that hunted mainly in open spaces, is indicative of the spread of savannah-like landscapes typical of the outskirts of the mammoth steppe. Changes in the species composition of bats occurring during the late Pleistocene and Holocene in southern Primorye adequately reflect the climate and landscape changes.

© 2016 Elsevier Ltd and INQUA. All rights reserved.

1. Introduction

The southern part of the Far Eastern region of Russia is characterized by increased diversity and the mixed nature of the fauna present, due to the peculiarities of geographical location, diversity of today's natural environment, and history of species expansion. The most significant changes in the distribution of animals relevant to the present occurred in the period of global climate changes in the late Pleistocene. There was a periodic offset of the area's fauna in the meridional direction on the eastern slopes of the Sikhotealin and throughout the entire Pleistocene refugium due to the humid climate (Krestov et al., 2009; Momohara et al., 2016). The unique features of that region may be elucidated by studying the history of faunal interchange directly in the local fossil record (Kim et al., 2015; Puzachenko et al., in press). The aim of this paper is to

describe fossil and subfossil records of bat fauna in the Medvezhyi Klyk cave (43° 01'43 "N, 133° 01'23"E) and, taking into account the environmental characteristics of particular species, to identify opportunities for the reconstruction of the paleogeographic condition and changes in the species composition of bats.

Of the 18 species of bats that presently live in the south of the Russian Far East (Tiunov, 2011), the majority are located on the southern or the northern limits. Fossil and subfossil findings of bats in the region are few. Most of these specimens are only defined to the genus (Ovodov, 1974, 1977). The most complete material was obtained from the Bliznetz cave location (Tiunov et al., 1992; Tiunov, 1997).

2. Materials and methods

The samples investigated in this study comprises subfossil and fossil teeth originating from the Medvezhyi Klyk cave deposits, situated in the central part of the Lozovyi mountain range (spurs of

E-mail address: tiunov@ibss.dvo.ru.

the mountain system of Sikhote-Alin, Primorsky Krai, Russia) (see Fig. 1). The entrance (2.0 × 0.55 m) is located on the crest of the watershed in the northern part of the ridge at an altitude of 465 m above sea level. The cave is a vertical cavity of karst origin that is 17.4 m deep. The bottom of the cave is shaped like an elongated oval, 5.3 m in length and 1.3 m in width. The area of the excavation was 1.0 × 0.5 m. The 13 lithological layers recognized in the Medvezhyi Klyk cave compose a total excavation depth of 5.3 m (Panasenko and Tiunov, 2010; Tiunov et al., 2016). Sediments were selected during excavation with a conditional horizon of 5–10 cm. The sediment samples taken in the field were screen-washed with 1.0 mm mesh size for microvertebrates. Location layers 1, 2, 3, 5, 7, 9, 11, 12, and 13 were horizontal, and fossil material was not mixed. The material of the layers (6, 7, 8, 9) with an inclined position (see Fig. 2) has been merged in the analysis. According to the radiocarbon dating of a humerus of a brown bear found in layer 7 (1.08–1.18 m), the estimated age of this layer is 13,790–14,200 years (GIN-13479) (Tiunov et al., 2016). Estimates for the other layers range from 7000 to 45,000 years BP. The accumulation of the bone remains of bats is probably the result of both natural mortality of animals during hibernation, and the result of birds of prey making their nests in the proximal part of the cave. The fossil remains do not show major morphological and/or morphometric differences from living species.

Bat remains are housed in the collection of the Institute of Biology and Soil Science, Russian Academy of Sciences in Vladivostok.

3. Results and discussion

There were 4715 fragments of the axial skull and lower jaws of 15 species of bats found. In analysing the results, only the lower jaw fragments of bats, which prevailed in the cave sediments, were considered. From 3544 lower jaw fragments, 2990 were identified to match a species (see Table 1).

The greatest number and the richest species composition of bat bone remains were found in the lower layer (13–11) deposits at depths ranging from 5.4 m to 2.53 m. According to preliminary estimates, the age of these layers corresponds to the marine oxygen isotope stage (MIS) 3. The fragments of skulls and lower jaws found were identified as belonging to the following species: *Rhinolophus nippon*, *Plecotus ognevi*, *Myotis rufoniger*, *M. petax*, *M. macrodactylus*, *M. bombinus*, *M. frater*, *M. gracilis*, *M. ikonnikovi*, *Hypsugo alashanicus*, *Eptesicus nilssonii*, *Murina hilgendorfi*, *Murina ussuriensis*, *Murina* sp.

Deposits accumulated in layers 9–10 probably date to the Last Glacial Maximum, approximately 18–20 ka BP. These layers revealed a minimal amount of bat bone remains belonging to *Plecotus ognevi* and *Murina hilgendorfi*. Currently, these two species are the most common hibernating in the caves south of the Primorsky Territory, even in small, relatively cool caves (Tiunov, 1985).

Species diversity of bat bone residues in more recent layers gradually increased, reaching a maximum in layer 5, formed during the Holocene optimum of approximately 7 ka BP (from 0.49 m to 0.95 m). In this layer, the fossil remains of *Plecotus ognevi*, *M. bombinus*, *M. frater*, *M. ikonnikovi*, *Hypsugo alashanicus*, *Eptesicus pachyomus*, and *Murina hilgendorfi* were discovered.

Presently, single individuals of *Murina hilgendorfi* and *Plecotus ognevi* are only occasionally found hibernating in caves.

Bone remains of *Murina hilgendorfi* were the most numerous and were found in all deposit layers. The relative amount of their bone depositions in a layer varied from 40 to 100% (see Fig. 3). In a layer, the mean relative abundance was 66%. It is obvious that this species existed at least 40 thousand years ago, and is now present in all of the Primorye territory (Tiunov, 1985), dominating and hibernating in caves. Considering that this is a typical forest species, it is clear that it has a continuous presence in forest landscapes, even in times of climatic maxima.

Plecotus ognevi is also a forest species. With regards to the number of fossil remains in depositions before Last Glacial




Fig. 1. Location of the Medvezhyi Klyk cave.

Download English Version:

<https://daneshyari.com/en/article/5114030>

Download Persian Version:

<https://daneshyari.com/article/5114030>

[Daneshyari.com](https://daneshyari.com)