



## CASE REPORT

# Mitral valve aneurysm: A serious complication of aortic valve endocarditis



Maria João Sousa\*, Vasco Alves, Sofia Cabral, Nuno Antunes,  
Luís Sousa Pereira, Filomena Oliveira, João Silveira, Severo Torres

*Serviço de Cardiologia, Centro Hospitalar do Porto, Porto, Portugal*

Received 18 November 2015; accepted 6 February 2016

Available online 17 October 2016

### KEYWORDS

Mitral valve;  
Aneurysm;  
Aortic valve;  
Endocarditis;  
Echocardiography

**Abstract** Mitral valve aneurysms are rare and occur most commonly in association with aortic valve endocarditis. Transesophageal echocardiography is the most sensitive imaging modality for the diagnosis of this entity and its potential complications, such as leaflet rupture and mitral regurgitation, which mandate prompt surgical intervention.

We present the case of a 70-year-old male patient with aortic valve endocarditis complicated with a ruptured aneurysm of the anterior mitral valve leaflet and associated severe mitral regurgitation, diagnosed by transesophageal echocardiography, with impressive images. We hypothesized that the aneurysm developed through direct extension of infection from the aortic valve or from a prolapsing aortic vegetation, with abscess formation and subsequent rupture and drainage. This case highlights the importance of appropriate imaging for early detection and timely surgical intervention (repair or replacement) to prevent fatal outcomes. © 2016 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. All rights reserved.

### PALAVRAS-CHAVE

Válvula mitral;  
Aneurisma;  
Válvula aórtica;  
Endocardite;  
Ecocardiografia

**Aneurisma da válvula mitral: uma complicação grave da endocardite valvular aórtica**

**Resumo** Os aneurismas da válvula mitral são raros e, habitualmente, encontram-se associados a endocardite da válvula aórtica. O ecocardiograma transesofágico é a modalidade imagiológica mais sensível para o diagnóstico desta entidade e das suas potenciais complicações, tais como perfuração e regurgitação mitral, que exigem intervenção cirúrgica urgente.

\* Corresponding author.

E-mail address: [maria.sousa.101@sapo.pt](mailto:maria.sousa.101@sapo.pt) (M.J. Sousa).

Apresentamos o caso clínico de um doente de 70 anos, do sexo masculino, com endocardite da válvula aórtica complicada por aneurisma perfurado do folheto anterior da válvula mitral e regurgitação mitral grave associada, diagnosticada por ecocardiograma transesofágico com imagens muito ilustrativas. Admitimos como hipótese, mais provável, que o aneurisma se desenvolveu por extensão direta da infeção a partir da válvula aórtica ou do prolapso de uma vegetação aórtica, com formação de abscesso e subsequente rotura e drenagem. Este caso realça a importância das modalidades imagiológicas adequadas no diagnóstico precoce do aneurisma da válvula mitral e da intervenção cirúrgica atempada (plastia ou substituição) na prevenção de eventos fatais.

© 2016 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L.U. Todos os direitos reservados.

## Case report

A 70-year-old male patient with type 2 diabetes mellitus and chronic hepatitis C virus infection presented to the emergency department with a four-month history of intermittent fever, night sweats, weight loss and progressive dyspnea. There was no past history of cardiovascular disease or illicit drug abuse. On physical examination, he was febrile (38.1 °C), his pulse was 124 bpm, blood pressure was 114/73 mmHg and he had signs of left heart failure. Cardiac auscultation disclosed a grade 3/6 pansystolic murmur at the apex. Blood cultures were positive for *Streptococcus viridans*.

Transthoracic echocardiography (TTE) showed multiple aortic valve vegetations (some prolapsing into the left ventricular outflow tract), mild aortic regurgitation and a large sacular bulge originating from the anterior mitral valve leaflet and protruding into the left atrium with severe mitral regurgitation (Figure 1); left ventricular systolic function was normal. With transesophageal echocardiography (TEE) it became clear that the sacular bulging on the mitral valve was an anterior mitral valve leaflet aneurysm with two rupture sites (Figure 2). Color flow Doppler showed two regurgitant jets: one through the coaptation margin of the mitral valve leaflets and the other across the perforated aneurysm (Figure 3). There was no evidence of either aortic ring abscess formation or extension along the mitral-aortic intervalvular fibrosa.

Antibiotic therapy was started with vancomycin plus gentamicin (due to penicillin allergy) for infective endocarditis (IE). After clinical stabilization, he was referred for urgent cardiac surgery. Given his clinical stability, surgery was delayed for four more days of antibiotic therapy. The patient died suddenly one day before scheduled surgical intervention.

## Discussion

Mitral valve aneurysms (MVAs) are uncommon but potentially serious complications, occurring most frequently in association with aortic valve endocarditis.<sup>1-4</sup> The first case was reported by Morand in 1729.<sup>5</sup> In 1970, Gonzalez-Lavin et al.<sup>6</sup> reported secondary involvement of the mitral

valve with leaflet aneurysm in two (3.4%) patients out of 58 who underwent aortic valve surgery for IE. Similar results were reported in 1992 by Karalis et al.<sup>7</sup> in a group of 55 patients with aortic valve endocarditis evaluated by TEE. More recent reports present a very low incidence, around 0.2-0.29% of patients who undergo TEE.<sup>2,4</sup>

Although most frequently associated with IE, other non-infectious etiologies have been reported including congenital cardiac structural defects, extreme forms of mitral valve prolapse, Libman-Sacks endocarditis and connective tissue disorders such as Marfan syndrome, Ehlers-Danlos syndrome, pseudoxanthoma elasticum and osteogenesis imperfecta.<sup>4,9-13</sup>

Because MVA is rare in the absence of IE, an infectious etiology will be at least partly responsible for leaflet degeneration. Proposed mechanisms for the development of MVA in the setting of aortic valve endocarditis include direct extension of the infection through the mitral-aortic intervalvular fibrosa to the anterior mitral valve leaflet with abscess formation and subsequent drainage, seeding of the anterior mitral valve leaflet by the aortic regurgitant jet ('jet lesion'),<sup>1,3,8,14-16</sup> or through direct contact with an aortic vegetation prolapsing into the left ventricular outflow tract ('kissing vegetation'), especially if the vegetation exceeds 6 mm in size.<sup>17</sup> These processes are thought to cause localized inflammation, weakness, dissection, and subsequent expansion of the valvular tissue, which in turn protrudes toward the left atrium due to ventricular systolic pressure, leading to aneurysm formation.<sup>2,18-21</sup> In our case, because aortic regurgitation was only mild, we believe that the most likely mechanism for aneurysm development was previous abscess formation (secondary to direct extension of infection from the aortic valve or a prolapsing aortic vegetation) followed by rupture and drainage.

The diagnosis may be suspected by TTE but TEE is more sensitive for detecting MVA, leaflet rupture and mitral regurgitation.<sup>2,4,22-25</sup> Recently, a few reports have highlighted the role of real-time three-dimensional echocardiography to better define the anatomical features of MVA preoperatively.<sup>16,26-28</sup>

Download English Version:

<https://daneshyari.com/en/article/5126571>

Download Persian Version:

<https://daneshyari.com/article/5126571>

[Daneshyari.com](https://daneshyari.com)