Accepted Manuscript

Title: Separation of enantiomers on chiral stationary phase based on cellulase: effect of preparation method and silica particle diameters on chiral recognition ability

Author: Hisami Matsunaga Jun Haginaka

PII: S0021-9673(16)30676-8

DOI: http://dx.doi.org/doi:10.1016/j.chroma.2016.05.069

Reference: CHROMA 357596

To appear in: Journal of Chromatography A

Received date: 25-3-2016 Revised date: 19-5-2016 Accepted date: 20-5-2016

Please cite this article as: Hisami Matsunaga, Jun Haginaka, Separation of enantiomers on chiral stationary phase based on cellulase: effect of preparation method and silica particle diameters on chiral recognition ability, Journal of Chromatography A http://dx.doi.org/10.1016/j.chroma.2016.05.069

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Separation of enantion	ners on chiral stationary phase based on cellulase:
effect of preparation n	nethod and silica particle diameters on chiral
recognition ability	

Hisami Matsunaga, Jun Haginaka*

School of Pharmacy and Pharmaceutical Sciences, Mukogawa Women's University, 11-68,

Koshien Kyuban-cho, Nishinomiya 663-8179, Japan

Corresponding Author. Tel: +81-798-45-9949; fax: +81-798-41-2792.

E-mail address: haginaka@mukogawa-u.ac.jp (J. Haginaka).

Download English Version:

https://daneshyari.com/en/article/5135617

Download Persian Version:

https://daneshyari.com/article/5135617

<u>Daneshyari.com</u>