



King Saud University
Arabian Journal of Chemistry

www.ksu.edu.sa
www.sciencedirect.com



ORIGINAL ARTICLE

Degradation of acid red 17 dye with ammonium persulphate in acidic solution using photoelectrocatalytic methods

M. Thabet, A.A. El-Zomrawy *

Department of Chemistry, Faculty of Science, Al-Azhar University, Cairo, Egypt

Received 13 January 2011; accepted 3 March 2011

KEYWORDS

Acid red 17;
Ammonium persulfate;
Degradation;
Photoelectrocatalytic

Abstract Advanced oxidation processes (AOPs) have proved very effective in the treatment of the various hazardous organic pollutants in water. The photoelectrocatalytic degradation of azo dye acid red 17 (AR-17) with ammonium persulfate (APS) was studied. Various operational parameters effect on the photoelectrocatalytic degradation rate, such as pH and the amount of APS oxidant, were investigated. The rate of degradation of dye with these composites followed pseudo-first order kinetics in the dye concentration. The results were observed and it was found that the dye decolorization was enhanced using pH 2.0. Also, the degradation rates were found to be strongly influenced by the increasing of the APS oxidant. The addition of $(\text{NH}_4)_2\text{S}_2\text{O}_8$ considerably increased the removal effectiveness due to the generation $\text{SO}_4^{\cdot-}$ radicals.

© 2011 King Saud University. Production and hosting by Elsevier B.V. All rights reserved.

1. Introduction

The introduction of waste products in the environment is a world wide problem that has been highlighted by various environmentalist groups. Dyes are widely used in industries such as textiles, rubber, paper, plastics, cosmetics, foods, etc., to color their products. The dyes are invariably left as the major waste

in these industries. Due to their chemical structures, dyes are resistant to fading on exposure to light, water and many chemicals and, therefore, are difficult to be decolorized once released into the aquatic environment (Sharma and Janveja, 2008). Many of the organic dyes are hazardous and may affect aquatic life and even the food chain (Sheikh and Rana, 2009). Release of these dyes in water stream is esthetically undesirable and has serious environmental impact. Due to intense color they reduce sunlight transmission into water hence affecting aquatic plants, which ultimately disturb aquatic ecosystem; in addition they are toxic to humans also.

The release and accumulation of dyes in suspension solution from tanneries, textile, paper and other industries formed in inland waters produce tremendous chemico-azo stress on aquatic organism including fishes and some time results in their mass mortality. The removal of dyes from industrial waste before they are discharged into the water bodies is therefore very important from health and hygiene point of view and for environmental protection.

* Corresponding author. Tel.: +20 106416041.
E-mail address: azomrawy@yahoo.com (A.A. El-Zomrawy).

1878-5352 © 2011 King Saud University. Production and hosting by Elsevier B.V. All rights reserved.


Peer review under responsibility of King Saud University.
doi:10.1016/j.arabjc.2011.03.001



Production and hosting by Elsevier

Methods for treating dye wastewaters consist of various chemical, physical and biological processes. These include: adsorption (Bilal, 2004), nanofiltration (Chakraborty et al., 2003; Capara et al., 2007), colloidal gas aphrons (Roy et al., 1992), electro coagulation (Alinsafi et al., 2004), coagulation and precipitation (Liu et al., 2003), advanced chemical oxidation (Arslan et al., 2000), electrochemical oxidation (Riera-Torres and Gutiérrez, 2010), photo-oxidation (Patricia et al., 2003), predispersed solvent extraction (Mahmoud et al., 2007), ozonation, supported liquid membrane (Mahmoud et al., 2007) and liquid-liquid extraction (Venkateswaran and Palanivelu, 2005). The reaction rate of azo dyes depends on the basic structure of the molecule and on the nature of auxiliary groups attached to the aromatic nuclei of dyes.

The persulphate also could be a good candidate for the photochemical oxidation technique. Persulphate ($S_2O_8^{2-}$) is a strong oxidant ($E_0 = 2.05$ V) which has been used widely in the petroleum industry for the treatment of hydraulic fluids or as a reaction initiator in the petrochemical industry (Daneshvar et al., 2007). The persulphate is normally available as a salt associated with ammonium, sodium, or potassium. It has also been reported to be effective for degrading organics in hazardous wastewaters in acidic or basic media through direct chemical oxidation (DCO), where persulphate is used as a sacrificial reagent (McCallum et al., 2000; Lau et al., 2007). The thermal or photochemical activated decomposition of $S_2O_8^{2-}$ ion to $SO_4^{\cdot-}$ radical has been proposed as a method to accelerate the process because the reactions of persulphate are generally slow at normal temperature (Lau et al., 2007). Photolysis of $S_2O_8^{2-}$ produces sulphate radical anions $SO_4^{\cdot-}$ with 100% photochemical efficiency, yielding $SO_4^{\cdot-}$ as summarized in the following reactions (Yu et al., 2004):




Similar to hydroxyl radicals, sulphate radicals may react with organics by electron transfer, hydrogen abstraction, or addition mechanisms (Anipsitakis et al., 2006).

In this work, the decoloration of acid red 17 by $S_2O_8^{2-}$ photochemically assisted and photoelectrochemically assisted in the work in different pH values has been studied. The study is based on monitoring reduction in the solution absorbance at the wavelength that corresponds to the maximum absorbance of the dye sample.

2. Materials and experimental methods

Bordeaux R, molecular formula = $C_{20}H_{12}N_2Na_2O_7S_2$, molecular weight = 502.42, $\lambda_{max} = 511 \pm 2$ nm, class = azo dye, C.I. number 16180, C.I. name acid red 17, and having structure:

Discoloration efficiency of AR-17 was measured by electrocatalytic, photocatalytic, and photoelectrocatalytic methods at pH 1.0 and 20 mA between two graphite electrodes. Initial



Chemical structure of acid red 17

concentrations of acid red, AR-17, and ammonium persulphate, APS, were 0.1 and 2.0 mM, respectively. Samples were withdrawn from a sample point at certain time intervals and analyzed for decoloration of acid red 17 dye. Thus, an optimum pH for maximum degradation of the aqueous AR-17 dye solutions by PECO method was studied. The reactions were carried out at different pH values in the range of 0.5–3.0 by adding H_2SO_4 or NaOH to adjust the pH value. PCO and PECO experiments of acid red dye (100 ml, 0.1 mmol) were carried out using UV Lamp (256 nm) of intensity 6 W.

Discoloration of dye solutions was checked and controlled by measuring the absorbance of dye at different intervals using UVmini-1240 SHMADZU, UV-vis spectrophotometer. The variation of pH of the sample before and after the irradiation was measured with a digital pH-meter, PHYWE 13702-93.

The degree of decoloration, i.e. the removal degree of color at the λ_{max} of the sample (511 nm) was calculated with the relation

$$\% = [(A_0 - A_t)/A_0] \times 100$$

where A_0 is the initial absorbance of the wastewater sample and A_t is the absorbance at the time t of the photodegradation.

Kinetic experiments were conducted by using PECO method. Kinetic decoloration behavior of AR-17 was investigated at different pH's. The effects of initial pH value on the degradation process were conducted by preparing 200 mL solutions containing 0.1, 2.0 mM of AR-17 and APS, respectively. On the other hand; the effects of initial APS concentrations were investigated at pH 2.0 and 0.1 mM of dye. All the samples at different time intervals (0–60 min) were taken.

3. Results and discussion

3.1. Decolorization efficiency

The results of decolorization of AR-17 dye with APS were obtained by the various methods; electrocatalytic oxidation (ECO), photocatalytic oxidation (PCO) and combination with each other, photoelectrocatalytic oxidation (PECO) in acidic medium. By using (PCO) the dyes molecules absorb photon, the UV energy will be transferred to the dyes molecules and the electronic structures of the dyes are excited. Thus, molecules excited states are unstable and short lived and will disperse their excitation energy in some physical way or undergo chemical reactions and form new species. The combination photoelectrocatalytic oxidation (PECO) was more effective in the decoloration of dye at pH 2.0 than (ECO) and (PCO) alone, where the AR-17 dye removal efficiency was 24.7% after 30 min, Fig. 1a and b.

The dye is attacked by UV photon and by hydroxyl and sulphate radicals, which are powerful oxidants generated from the electro and the photolysis of H_2O and ammonium persulphate by UV light, making possible the degradation of the dyes. The destruction of azo bond $-N=N-$ in the chromophore of azo dyes led to the decoloration of the dye solutions.

The effects of PECO method on the decoloration efficiency of AR-17 at various pH (0.5, 1.0, 2.0, 3.0) and in presence of 2.0 mM of APS were investigated. Discoloration of AR-17 dye at pH 2.0 was more efficient than other pH values under this study, Fig. 2, suggesting that the pH value would affect the amount of OH^{\cdot} and $SO_4^{\cdot-}$ generation, and the preferable condition for OH^{\cdot} and $SO_4^{\cdot-}$ generation was under acidic conditions.

Download English Version:

<https://daneshyari.com/en/article/5142480>

Download Persian Version:

<https://daneshyari.com/article/5142480>

[Daneshyari.com](https://daneshyari.com)