

Accepted Manuscript

Synthesis and molecular modelling studies of phenyl linked oxadiazole-phenylhydrazone hybrids as potent antileishmanial agents

Muhammad Taha, Nor Hadiani Ismail, Syahrul Imran, El Hassane Anouar, Manikandan Selvaraj, Waqas Jamil, Muhammad Ali, Syed Muhammad Kashif, Fazal Rahim, Khalid Mohammed Khan, Mohd Ilham Adenan

PII: S0223-5234(16)31025-X

DOI: [10.1016/j.ejmech.2016.12.019](https://doi.org/10.1016/j.ejmech.2016.12.019)

Reference: EJMECH 9110

To appear in: *European Journal of Medicinal Chemistry*

Received Date: 31 May 2016

Revised Date: 6 December 2016

Accepted Date: 9 December 2016

Please cite this article as: M. Taha, N.H. Ismail, S. Imran, E.H. Anouar, M. Selvaraj, W. Jamil, M. Ali, S.M. Kashif, F. Rahim, K.M. Khan, M.I. Adenan, Synthesis and molecular modelling studies of phenyl linked oxadiazole-phenylhydrazone hybrids as potent antileishmanial agents, *European Journal of Medicinal Chemistry* (2017), doi: 10.1016/j.ejmech.2016.12.019.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.




Graphical Abstract

**Synthesis and Molecular Modelling
Studies of Phenyl Linked
Oxadiazole-Phenylhydrazone Hybrids as Potent
Antileishmanial Agents**

Muhammad Taha^{*a,b}, Nor Hadiani Ismail^{a,b}, Syahrul Imran^{a,b}, El Hassane Anouar^c, Manikandan Selvaraj^{d,e}, Waqas Jamil^f, Muhammad Ali^f Syed Muhammad Kashif^f, Fazal Rahim^g, Khalid Mohammed Khan^h, Mohd Ilham Adenan^{a,b}

^aAtta-ur-Rahman Institute for Natural Product Discovery, Universiti Teknologi MARA (UiTM), Puncak Alam Campus, 42300 Bandar Puncak Alam, Selangor, Malaysia; ^bFaculty of Applied Science UiTM, 40450 Shah Alam, Selangor, Malaysia; ^cChemistry Department, College of Sciences and Humanities, Prince Sattam bin Abdulaziz University, Suadia Arabia; ^dIntegrative Pharmacogenomics Institute (iPROMISE), Universiti Teknologi MARA (UiTM), Puncak Alam Campus, 42300 Bandar Puncak Alam, Selangor Darul Ehsan, Malaysia; ^eFaculty of Pharmacy, Universiti Teknologi MARA (UiTM), Puncak Alam Campus, 42300 Bandar Puncak Alam, Selangor Darul Ehsan, Malaysia; ^fInstitute of Advance Research Studies in Chemical Sciences, University of Sindh Jamshoro, Hyderabad 76080, Pakistan; ^gDepartment of Chemistry, Hazara University, Mansehra, Pakistan; ^hH. E. J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi-75270, Pakistan



Compound **10** showed $0.95 \pm 0.01 \mu\text{M}$ as compare to standard drug pentamidine ($\text{IC}_{50} = 7.02 \pm 0.09 \mu\text{M}$)

Download English Version:

<https://daneshyari.com/en/article/5158885>

Download Persian Version:

<https://daneshyari.com/article/5158885>

[Daneshyari.com](https://daneshyari.com)