Accepted Manuscript

A multi-technique approach to assess the fate of high-temperature biochar in soil and to quantify its effect on soil organic matter composition

Lydia Paetsch, Carsten W. Mueller, Cornelia Rumpel, Š árka Angst, Alexandra C. Wiesheu, Cyril Girardin, Natalia P. Ivleva, Reinhard Niessner, Ingrid Kögel-Knabner

PII: S0146-6380(17)30039-6
DOI: http://dx.doi.org/10.1016/j.orggeochem.2017.06.012
Reference: OG 3568

To appear in: *Organic Geochemistry*

Received Date: 19 January 2017
Revised Date: 8 June 2017
Accepted Date: 9 June 2017

Please cite this article as: Paetsch, L., Mueller, C.W., Rumpel, C., Angst, S., Wiesheu, A.C., Girardin, C., Ivleva, N.P., Niessner, R., Kögel-Knabner, I., A multi-technique approach to assess the fate of high-temperature biochar in soil and to quantify its effect on soil organic matter composition, *Organic Geochemistry* (2017), doi: http://dx.doi.org/10.1016/j.orggeochem.2017.06.012

A multi-technique approach to assess the fate of high-temperature biochar in soil and to quantify its effect on soil organic matter composition.

Lydia Paetsch[*1], Carsten W. Mueller[1], Cornelia Rumpel[2,3], Šárka Angst[4], Alexandra C. Wiesheu[5], Cyril Girardin[2], Natalia P. Ivleva[5], Reinhard Niessner[5,6], Ingrid Kögel-Knabner[1,7]*

[1]Chair of Soil Science, Technical University of Munich, Emil-Ramann-Strasse 2, 85354 Freising, Germany

[2]INRA, ECOSYS, UMR INRA-AgroParisTech, Thiverval-Grignon, CentreAgroParisTech, bâtiment EGER, 78850, France

[3]CNRS, IEES, UMR UPMC-CNRS-UPEC-INRA-IRD, CentreAgroParisTech, bâtiment EGER, Thiverval-Grignon, 78850, France

[4]Department of Ecosystem Biology, Faculty of Science, University of South Bohemia, Branišovská 31, 370 05 České Budějovice, Czech Republic

[5]Institute of Hydrochemistry, Chair for Analytical Chemistry, Technical University of Munich, Marchioninistrasse 17, 81377 Munich, Germany