

## CHARACTERISTICS OF TWO GLUTAMINE SYNTHETASE ISOZYMES IN SOYBEAN

SEONG-MO KANG and THEODORE HYMOWITZ

Department of Agronomy, University of Illinois, Urbana, IL 61801, U.S.A.

(Revised received 21 October 1987)

**Key Word Index**—*Glycine max*; Leguminosae; soybean; L-glutamate ammonia ligase; glutamine synthetase.

**Abstract**—Soybean plants [*Glycine max* (L.) Merr. cv. Williams] contained two chromatographically distinct isoforms of glutamine synthetase, GS-1 and GS-2. GS-1 was localized in the soluble fraction while GS-2 was in the chloroplasts. GS-2 occurred only in isolated chloroplasts, and etiolation or cycloheximide treatment of light-grown plants resulted in a dramatic decrease of GS-2 activity. Although there were some minor differences in their characteristics, the isoenzymes did not differ significantly. The increase in total activity during the early phases of growth was due largely to the increase in GS-2 activity. The activity in dark-grown plants, when exposed to light, reached almost to the level of light-grown plants within 48 hr. When the two isoenzymes were separated by DEAE-Sephacel chromatography, GS-1 accounted for 20% of total activity recovered. Although this proportion was found in both four- and 16-day-old seedlings, this conclusion may not be extrapolated to intact seedlings because of the activity loss during purification and different stability of the isoenzymes. On a whole seedling basis, the activity of the enzyme increased during the three-week period of germination and early growth. The reverse was true on a unit weight basis. Specific activities of roots and primary stems were much higher than any other parts of the plant.

### INTRODUCTION

Glutamine is the first form of organic nitrogen fixed from atmospheric nitrogen. The importance of glutamine has also been indicated by its role as a nitrogen donor for the synthesis of various metabolites. Since glutamine synthesis is catalysed by glutamine synthetase (EC 6.3.1.2), the importance of the enzyme in determining the overall nitrogen economy of the plants is self-evident. In addition to its role in nitrogen assimilation, the enzyme has also been implicated in the recycling of ammonia released during photorespiration [1,2].

Although glutamine synthetase (GS) has been studied in a number of plant species for some time, only recently it has been shown to exist as at least two different isoenzymes designated as GS-1 and GS-2 in a limited number of plant species [3,4]. Studies on cellular localization of the isoenzymes have indicated that GS-1 is present in the cytoplasm while GS-2 is present in the chloroplasts.

In this paper, we report the presence and cellular localization of two chromatographically distinct isoenzymes of GS in soybean. Special attention was given to the relative proportion of GS-1 and GS-2 in the whole plant, and development of their activity during the early phases of growth. Unlike most studies which have dealt with a particular tissue of a plant, mostly leaves, we believe that it is important to understand GS in the whole plant.

### RESULTS

#### Enzyme purification

Table 1 shows the results from an enzyme purification. Although a 25% loss of activity resulted, ammonium

sulphate precipitation of the enzymes was useful in removing the bulk of proteins in soybean seeds and seedlings.

Separation of GS into two isoenzymes was easily achieved at the DEAE-Sephacel step (Fig. 1A) at the expense of ca 60% loss of activity from the crude extract. They were eluted from the column at 0.16 and 0.27 M NaCl respectively. The activity peak fractions of GS-1 and GS-2 were rechromatographed individually on a hydroxyapatite column. GS-1 and GS-2 were eluted from the column at 0.065 M and 0.165 M phosphate respectively (Fig. 1B and C). Recovery was 4 and 9% for GS-1 and GS-2 and they were purified 94- and 143-fold respectively (Table 1). GS-1 recovery of the DEAE-Sephacel fraction from the hydroxyapatite step was 55%; the recovery of GS-2 activity was only 26%.

#### Characteristics of the two isoenzymes

The pH optima of GS-1 activity were 7.5 and 5.5 in the presence of  $Mg^{2+}$  and  $Mn^{2+}$ , respectively. Those of GS-2 were 0.5 pH unit higher than GS-1. There was a sharp decline in the  $Mg^{2+}$ -dependent activity of both isoenzymes below pH 5.0, while the  $Mn^{2+}$ -dependent activity declined sharply above pH 8.0. The  $Mn^{2+}$ -dependent activity was 32 to 34% of the  $Mg^{2+}$ -dependent activity with both isoenzymes.

There were no significant differences between GS-1 and GS-2 in their apparent  $K_m$  values for various substrates. The values of GS-1 and GS-2 respectively were in mM: glutamate, 7.56 and 6.45; ATP, 0.94 and 0.87;  $NH_2OH$ , 0.48 and 0.52;  $NH_4^+$ , 0.038 and 0.029;  $Mg^{2+}$ , 13.14 and 14.47;  $Mn^{2+}$ , 0.77 and 1.15.

Both GS-1 and GS-2 showed maximum activity at 42°, with a sharp decline in their activity above this temperature. GS-1 and GS-2 retained only 27 and 7% of their




Fig. 1. Elution profiles of two glutamine synthetase isoenzymes on a DEAE-Sephacel (A) and hydroxyapatite column (B, C). The two peak fractions of I and II in A were rechromatographed as shown in B and C, respectively. Proteins were eluted with a linear gradient of NaCl for A and K-Pi for B and C, with 5 ml (A) and 1.5 ml fractions (B and C). Every other fraction was assayed for enzyme activity.

activity, respectively, when they were preincubated at 55° for 10 min. When ATP was added in the preincubating enzyme solutions, both enzymes were quite stable up to 50°. At 55°, the residual activities of GS-1 and GS-2 were 72 and 51% respectively.

#### Localization of GS-2 in chloroplasts

GS activity of isolated chloroplasts was eluted from a DEAE-Sephacel column with 0.27 M NaCl, which corresponds to the same strength of NaCl where GS-2 was eluted in Fig. 1A. We therefore concluded that GS-2 in Fig. 1A originated from the chloroplasts. The treatment of light-grown plants with cycloheximide decreased GS-2

activity by 30% when the enzyme was extracted from the isolated chloroplasts. We routinely used 20 g of leaves to isolate chloroplasts and the enzyme therefrom, obtaining 56 units of activity in 5 mg protein. With the cycloheximide treatment, the recovery was 39 units in 3.8 mg protein.

A typical 16-day-old seedling contained *ca* 111 units of total activity. When the enzyme extract was separated by DEAE-Sephacel column chromatography, the recovered activities of GS-1 and GS-2 were 11.3 and 36.7 units. Thus, more than three-quarters of the total activity was from GS-2. When the seedlings were etiolated for six days before harvest, the activities of GS-1 and GS-2 recovered from the DEAE-Sephacel were 9.8 and 19.8 units respectively. Total activity declined 38% following etiolation; the decreases in activity of GS-1 and GS-2 were 13 and 46% respectively.

#### Activity changes during the early stage of growth

Total enzyme activity increased linearly during the first four-day germination in the light, and thereafter declined until day six (Fig. 2), after which per plant activity again gradually increased (Table 2). By contrast, in dark-grown seedlings, the activity increased to a maximum by 36 hr and then declined during the later sampling period. When dark-germinated seedlings were exposed to light, the activity increased sharply and, within 48 hr, the activity was similar to that of light-germinated seedlings. The activity of two-day-old dark-germinated seedlings, when exposed to light on day two, increased to 94% of the level of light-germinated seedlings by day four.

#### Distribution of enzyme activity in a whole seedling

Table 2 shows the changes in total activity in various parts of a seedling. In mature seeds, seed coats contained over three times more activity than the rest of the seed on a protein basis, although per seed activity in seed coat was just 4% of total activity. Total activity of cotyledon pairs declined dramatically during the first 10 days, after which it became more or less stabilized at 15–20% of a whole seedling. Specific activity of cotyledon pairs increased sharply in three weeks with some six-fold decline in proteins.

As plant mass increased, total activity in leaves and growing axis also increased; it accounted for 47% of a whole seedling grown for three weeks. However, specific activity of leaves and growing axis declined, suggesting that the rate of protein increase far exceeded that of activity increase during the same period. The relative proportion of root activity was stable at *ca* 20% level in all samples, so that the top/root ratio of total activity was four at the end of experiment. Total activity presented in Table 2 is the sum of the activity measured individually in various parts of a seedling. When the activity was measured from the extracts of whole seedlings without dividing them into various parts, total activity was 3–5% higher than the values presented in Table 2.

#### DISCUSSION

Soybean seedlings grown in the light contained two chromatographically distinct isoenzymes of GS (Fig. 1). When the two isoenzymes present in four-day-old, light-

Download English Version:

<https://daneshyari.com/en/article/5169982>

Download Persian Version:

<https://daneshyari.com/article/5169982>

[Daneshyari.com](https://daneshyari.com)