

Accepted Manuscript

New functionalized 8-hydroxyquinoline-5-sulfonic acid mesoporous silica (HQS-SBA-15) as an efficient catalyst for the synthesis of 2-thiohydantoin derivatives

Vaezeh Fathi Vavsari, Ghodsi Mohammadi Ziarani, Saeed Balalaie, Arezoo Latifi, Mehdi Karimi, Alireza Badiei



PII: S0040-4020(16)30657-3

DOI: [10.1016/j.tet.2016.07.034](https://doi.org/10.1016/j.tet.2016.07.034)

Reference: TET 27930

To appear in: *Tetrahedron*

Received Date: 12 April 2016

Revised Date: 3 July 2016

Accepted Date: 7 July 2016


Please cite this article as: Vavsari VF, Ziarani GM, Balalaie S, Latifi A, Karimi M, Badiei A, New functionalized 8-hydroxyquinoline-5-sulfonic acid mesoporous silica (HQS-SBA-15) as an efficient catalyst for the synthesis of 2-thiohydantoin derivatives, *Tetrahedron* (2016), doi: 10.1016/j.tet.2016.07.034.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Graphical abstract

New functionalized 8-hydroxyquinoline-5-sulfonic acid mesoporous silica (HQS-SBA-15) as an efficient catalyst for the synthesis of 2-thiohydantoin derivatives

Vaezeh Fathi Vavsari, Ghodsi Mohammadi Ziarani, Saeed Balalaie*, Arezoo Latifi, Mehdi Karimi, Alireza Badiei



Download English Version:

<https://daneshyari.com/en/article/5213288>

Download Persian Version:

<https://daneshyari.com/article/5213288>

[Daneshyari.com](https://daneshyari.com)