



Conversion of aldoximes into nitriles catalyzed by simple transition metal salt of the fourth period in acetonitrile



Xiao-Yun Ma, Ying He, Ting-Ting Lu, Ming Lu*

Chemical Engineering College, Nanjing University of Science and Technology, Nanjing 210094, China

ARTICLE INFO

Article history:

Received 21 October 2012

Received in revised form 13 January 2013

Accepted 21 January 2013

Available online 1 February 2013

Keywords:

Aldoxime

Nitrile

Amide

Acetonitrile

Transition metal salt

ABSTRACT

Conversion of aldoximes into nitriles catalyzed by simple transition metal catalysts, such as copper salts, nickel salts, cobalt salts, zinc salts, iron salts, and manganese salts in acetonitrile was investigated. All the metal salts display catalytic property in the conversion of aldoximes into nitriles and cupric acetate exhibits much higher activity than other catalysts. The corresponding amide was detected in almost all cases and acetonitrile was found to be involved in the conversion of aldoximes into nitriles.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Dehydration of aldoximes is a useful method for the synthesis of nitriles, which are important synthetic intermediates for pharmaceuticals, agricultural chemicals, dyes, and material sciences.¹ In the past 40 years, a variety of reagents, such as acid anhydrides,² acids,³ bases,⁴ acid chlorides,⁵ *N*-triflylimidazole,⁶ bromodimethylsulfonium bromide,⁷ phosphorus-containing compounds,⁸ etc. have been introduced for the dehydration of aldoximes to nitriles. However, the approach suffers from some disadvantages, such as the use of stoichiometric amounts of reagents and limitations arising from the sensitivity of some functional groups to the reaction conditions. Thus, the development of metal-catalyzed dehydration of aldoxime has received much attention. A number of metal catalysts, such as nickel catalysts,⁹ metal triflates,¹⁰ [RuCl₂(pcymene)],¹¹ [(CH₃CH₂CN)₂PtCl₄],¹² copper salts,¹³ zinc salts,¹⁴ Pd(OAc)₂,¹⁵ [(Ipr)AuCl]/AgBF₄,¹⁶ iron porphyrin,¹⁷ and tungsten-tin mixed hydroxide¹⁸ have been reported. In addition, supported catalyst¹⁹ and ionic liquid²⁰ were employed in the conversion of aldoximes into nitriles.

In our previous study,²¹ we disclosed an efficient method for the conversion of nitriles into the corresponding amides by employing acetaldoxime and simple transition metal catalysts of the fourth period. In the continuation of the study on simple transition metal


catalysts for the chemical transformation, we found that aldoximes were converted into the corresponding nitriles catalyzed by simple transition metal catalysts, such as copper salts, nickel salts, cobalt salts, zinc salts, iron salts, and manganese salts in acetonitrile. The corresponding amide was detected as a by-product in almost all cases and the yield of amide depends on the hydration of the nitrile product.

At the same time, we have noticed that the conversion of aldoximes into nitriles catalyzed by cupric acetate (Cu(OAc)₂),^{13a} CuCl₂ under ultrasound conditions^{13b} and nickel salts^{9b} in acetonitrile have been reported. In these literature, catalytic dehydration of aldoximes were performed efficiently with catalyst in acetonitrile. However, they did not mention the formation of amide except *p*-nitrobenzamide.^{9b} Moreover, they did not find that acetonitrile plays an important role in the reaction. In the course of our study, we are certain that the corresponding amide was obtained as a by-product in almost all cases and acetonitrile play dual roles as solvent and reagent in the reaction.

2. Results and discussion

In this paper, *p*-methoxybenzaloxime was selected as a model substrate to investigate the catalytic activity of various metal salts of the fourth period. As shown in Table 1, we found that all the metal salts displayed catalytic properties in the conversion of *p*-methoxybenzaloxime into *p*-methoxybenzoxime (Table 1, entries 2–14) and cupric acetate showed the highest catalytic activity (Table 1, entry 4). Both the cation and anion of the metal salt had an

* Corresponding author. Tel.: +86 025 84315514; fax: +86 025 84315030; e-mail address: lumingnjt@163.com (M. Lu).


Table 1
Screen of reaction conditions^a


Entry	Conditions	(Yield) (%) ^b	
		Nitrile	Amide
1	CH ₃ CN, reflux, 24 h.	—	—
2	CuO (0.1 equiv), CH ₃ CN, reflux, 24 h.	85	6
3	CuCl ₂ ·4H ₂ O (0.1 equiv), CH ₃ CN, reflux, 3 h	91	9
4	Cu(OAc) ₂ (0.1 equiv), CH ₃ CN, reflux, 0.25 h	96 (90)	4
5	Cu(OAc) ₂ (0.05 equiv), CH ₃ CN, reflux, 1.5 h	96	4
6	Ni(OAc) ₂ ·4H ₂ O (0.1 equiv), CH ₃ CN, reflux, 4 h	92	8
7	NiCl ₂ ·6H ₂ O (0.1 equiv), CH ₃ CN, reflux, 24 h	86	13
8	CoCl ₂ ·6H ₂ O (0.1 equiv), CH ₃ CN, reflux, 24 h	91	8
9	Co(OAc) ₂ ·4H ₂ O (0.1 equiv), CH ₃ CN, reflux, 12 h	93	7
10	ZnCl ₂ (0.1 equiv), CH ₃ CN, reflux, 24 h	92	7
11	Zn(OAc) ₂ ·4H ₂ O (0.1 equiv), CH ₃ CN, reflux, 24 h	93	6
12	FeCl ₂ ·4H ₂ O (0.1 equiv), CH ₃ CN, reflux, 24 h	13	Trace
13	MnCl ₂ ·4H ₂ O (0.1 equiv), CH ₃ CN, reflux, 24 h	12	Trace
14	Mn(OAc) ₂ ·4H ₂ O (0.1 equiv), CH ₃ CN, reflux, 24 h	16	Trace
15	Cu(OAc) ₂ (0.1 equiv), CH ₃ CN, rt, 24 h	91	Trace
16	Cu(OAc) ₂ (0.1 equiv), CH ₃ OH, rt, 24 h	—	—
17	Cu(OAc) ₂ (0.1 equiv), (CH ₃) ₂ CHOH, rt, 24 h	—	—
18	Cu(OAc) ₂ (0.1 equiv), toluene, rt, 24 h	—	—
19	Cu(OAc) ₂ (0.1 equiv), dioxane, rt, 24 h	—	—

^a Reaction conditions: *p*-methoxybenzaldehyde (2 mmol), solvent (5 mL).^b Determined by GC. In brackets the isolated yield is stated.

effect on the catalytic property. Treatment of *p*-methoxybenzaldehyde oxime with cupric acetate (Cu(OAc)₂), copper(II) chloride dihydrate (CuCl₂·2H₂O), copper(II) chloride tetrahydrate (CuCl₂·4H₂O), nickel(II) acetate tetrahydrate (Ni(OAc)₂·4H₂O) or cobalt(II) acetate (Co(OAc)₂·4H₂O) at refluxing temperature in anhydrous acetonitrile afforded 100% conversion (Table 1, entries 3–6 and 8). When we reduced the amounts of catalyst to 5 mol % (Table 1, entry 5), *p*-methoxybenzaldehyde oxime was converted to the corresponding nitrile and amide with complete conversion after a relatively long reaction time of 1.5 h. *p*-Methoxybenzamide was detected in each case and the yield of *p*-methoxybenzamide depends on the catalyst. When nickel(II) chloride hexahydrate (NiCl₂·6H₂O) was used as a catalyst, *p*-methoxybenzamide was obtained in relatively high yield (Table 1, entry 7). When iron salt or manganese salt was used as a catalyst, a trace amount of *p*-methoxybenzamide was detected (Table 1, entries 12–14). In addition, we found that the conversion of *p*-methoxybenzaldehyde oxime into *p*-methoxybenzitrile can proceed at ambient temperature when Cu(OAc)₂ was used as a catalyst (Table 1, entry 15). However, use of other solvent in place of acetonitrile resulted in no product when the reaction was carried out at room temperature (Table 1, entries 16–19). Therefore, we thought that acetonitrile may be involved in the reaction.

On the basis of the above results, we can conclude that using cupric acetate (Cu(OAc)₂) in acetonitrile at refluxing temperature is the optimal conditions for the conversion of aldoximes into nitriles. This protocol was subsequently applied to various aromatic aldoximes, aliphatic aldoxime and heterocyclic aldoximes. As shown in Table 2, various aldoximes including aromatic aldoximes (Table 2, entries 1–8), aliphatic aldoxime (Table 2, entry 9) and heterocyclic aldoximes (Table 2, entries 10–12) were converted into the corresponding nitriles and amides with complete conversion. Substrates bearing an electron-donating group proceed more effectively than aldoximes with an electron-withdrawing group (Table 2, entries 3 and 7), the sterically unhindered aldoximes were easier to convert into the corresponding nitriles than the sterically hindered aldoximes (Table 2, entries 2–4). Heterocyclic aldoximes with a heteroatom lone pair positioned *ortho* to the oximido group

Table 2
Conversion of various aldoximes using cupric acetate and acetonitrile^a


Entry	Substrate (R)	Time (h)	Yield (%) ^b	
			Nitrile	Amide
1	Ph	1	99 (91)	1
2	2-Cl-C ₆ H ₄	1.5	100 (95)	—
3	4-Cl-C ₆ H ₄	1	90 (83)	10
4	2-Cl-4-Cl-C ₆ H ₃	2	99 (93)	Trace
5	4-NO ₂ -C ₆ H ₄	1	88 (80)	12
6	3-MeO-4-MeO-5-MeO-C ₆ H ₂	0.75	95 (87)	5
7	4-OH-C ₆ H ₄	0.25	96 (89)	4
8	4-N(CH ₃) ₂ -C ₆ H ₄	0.25	99 (92)	Trace
9	n-C ₁₁ H ₂₃	1	99 (86)	Trace
10	4-Pyridyl	1.5	85 (77)	15
11	2-Furyl	0.25	57 (50)	43
12	2-Thienyl	0.25	69 (61)	31

^a Reaction conditions: aldoxime (2 mmol), Cu(OAc)₂ (0.2 mmol), anhydrous acetonitrile (5 mL), reflux.^b Determined by GC. In brackets the isolated yield is stated.

show high reactivity and these aldoximes were converted to the corresponding nitriles and amides with complete conversion after a short reaction time of 15 min (Table 2, entries 11–12). From Table 2, it can be seen that the corresponding amide was obtained as a by-product in all cases except 2-chlorobenzaldehyde oxime. More amide was obtained in the case of aldoxime having electron-withdrawing group and aldoximes without steric hindrance. In the case of heterocyclic aldoxime with a heteroatom lone pair positioned *ortho* to the oximido group, the yield of the corresponding amide was reached over 30% (Table 2, entries 11–12). To our surprise, the above phenomenon of formation of amide is similar to the hydration of nitriles catalyzed by transition metal salt with the aid of acetaldoxime.²¹ Therefore, we guessed that the conversion of aldoximes to nitriles may not proceed via the routes reported in the literature^{9b,13a} and acetonitrile may be involved in the reaction.

To prove our guess, we selected (*E*)-furan-2-carbaldehyde oxime as a model substrate to investigate the reaction pathway. As can be seen from Table 2, furan-2-carbaldehyde oxime showed high reactivity and a complete conversion of furan-2-carbaldehyde oxime took place within 15 min at refluxing temperature. In fact, this reaction can proceed at ambient temperature and 100% of (*E*)-furan-2-carbaldehyde oxime was transformed into furan-2-carbonitrile and furan-2-carboxamide. As shown in Table 3, we found that (*E*)-furan-2-carbaldehyde oxime was not converted to furan-2-carbonitrile by the action of cupric acetate (Cu(OAc)₂) alone (Table 3, entry 1). When 0.1 equiv of acetonitrile was added, 100% of (*E*)-furan-2-carbaldehyde oxime was transformed into furan-2-carboxamide at room temperature (Table 3, entry 2). Therefore, we can conclude that acetonitrile was not only involved in the reaction but that it also regenerated in this reaction. To our surprise, a mixture of furan-2-carbonitrile and furan-2-carboxamide was obtained when we increased the amounts of acetonitrile (Table 3, entry 3) and the furan-2-carbonitrile was completely converted to furan-2-carboxamide when an additional acetaldoxime was added. Based on the above results, we proposed a possible reaction pathway as follows: (*E*)-furan-2-carbaldehyde oxime reacted with acetonitrile in the presence of transition metal catalyst to give furan-2-carbonitrile and acetaldoxime, and then furan-2-carbonitrile reacted with acetaldoxime in the presence of transition metal catalyst to afford furan-2-carboxamide and acetonitrile.^{21,22} According to the proposed reaction pathway, we know that the existence of massive acetonitrile impedes the hydration of nitrile on the basis of the principles of chemical equilibrium. Moreover, acetonitrile can also be hydrated in presence of

Download English Version:

<https://daneshyari.com/en/article/5218189>

Download Persian Version:

<https://daneshyari.com/article/5218189>

[Daneshyari.com](https://daneshyari.com)