



Chiral pool based synthesis of pyrrolidinium ionic liquids



Alexandra Brown^a, Victoria Hogan^b, Jacob Perry^a, Renuka Manchanayakage^{a,*}

^a Department of Chemistry, St. John Fisher College, 3690 East Avenue, Rochester, NY 14618, United States

^b Department of Chemistry, Susquehanna University, 514 University Avenue, Selinsgrove, PA 17870, United States

ARTICLE INFO

Article history:

Received 10 January 2017

Revised 24 January 2017

Accepted 30 January 2017

Available online 3 February 2017

Keywords:

Ionic liquids

Chirality

Pyrrolidine

Chiral solvents

Enantiomerically pure synthesis

ABSTRACT

Chiral ionic liquids show promising applications in various different fields. A series of pyrrolidinium-based chiral ionic liquids bearing a chiral cation, a chiral anion or both was prepared in good yields using an efficient, economic and simple pathway. The chirality was introduced using (L)-lactate and (L)-menthol derivatives. The resultant chiral compounds were characterized by both spectroscopy and polarimetry. We envision that these new chiral compounds can serve as effective reaction media and chiral catalysts for asymmetric reactions, which are presently being investigated in our lab.

© 2017 Elsevier Ltd. All rights reserved.

In the recent years, there has been a growing demand in the use of the room-temperature ionic liquids (RTIL) as new reaction media.¹ One of the main expected applications of ionic liquids is to replace the volatile organic solvents traditionally used in large scale in industry.² Ionic liquids possess a number of interesting properties over commonly used organic solvents, such as lack of significant vapor pressure, absence of flammability, tolerance for large temperature variations and ease of reuse.³ They have been extensively used in chemical syntheses and electrosyntheses.⁴

Chiral ionic liquids (CILs) are one of the latest branches of ionic liquid research and are promising chiral solvents for many applications.⁵ Chiral ionic liquids are particularly attractive due to their potential for chiral discrimination, as in asymmetric synthesis and optical resolution of racemates.⁶ Because of their highly-organized nature and ionic properties, an obvious potential exists for solvent-solute interaction that may provide a mechanism for substantial improvement over conventional chiral solvents.⁷ So far, chiral ionic liquids have been successfully applied in asymmetric synthesis, stereoselective polymerization, chiral stationary phases in chromatographic techniques, and chiral shift reagents in NMR spectroscopy.⁸ Other advantages of using chiral ionic liquids include the following: (1) the chiral ionic liquids are easily and inexpensively prepared and can be recycled; (2) the opposite enantiomers of the chiral ionic liquids can be produced in order to enantioselectively create the desired enantiomer in excess; (3) it is possible and easy to remove the chiral ionic liquids from the final

reaction mixture so that no interference is occurred. Therefore, it is important to have a range of chiral ionic liquids with different properties that can be used in various applications.

The more efficient, economic and simple way to prepare enantiomerically pure ionic liquids with central chirality is the use of precursors derived from the chiral pool either for the generation of the CIL's anion or cation or for both.⁹ Chirality has been introduced to the cation using various methods and the imidazolium cation has been widely used in these preparations.¹⁰ Under most circumstances, the imidazole-derived ionic liquids are considered to be stable and "inert" solvents, but the 2-position can be deprotonated to form a stabilized carbene which could be problematic for some applications.¹¹ This situation can be avoided by the use of pyrrolidinium and benzimidazolium cations. However, less attention has been given to the cations such as pyrrolidinium and benzimidazolium in the preparation of chiral ionic liquids. Additionally, recent research shows that some pyrrolidinium-based ionic liquids have excellent electrochemical and mechanical performances and are promising electrolytes for lithium batteries.¹²


Armed with this information, we discuss the synthesis and characterization of a series of pyrrolidinium-based chiral ionic liquids. Chirality is introduced to the cation by using a menthol derivative and to the anion by using a lactate salt. The presence of functional groups makes them potential task-specific chiral ionic liquids.¹³ The concept of tailor-made task-specific chiral ionic liquids is rather new, and the synthesis of a series of new chiral ionic liquids with unique properties will benefit the collection of compounds that is currently limited in scope.

* Corresponding author.

E-mail address: rmanchanayakage@sjfc.edu (R. Manchanayakage).

This project focused on the synthesis and characterization of pyrrolidinium-based chiral ionic liquids bearing a chiral cation, chiral anion or both using an economic, simple and efficient pathway. The enantiomerically pure ionic liquids can be prepared using chiral precursors such as aminoacids, aminoalcohols, hydroxyacids, amines, alkaloids or halogenoalkanes derived from a chiral pool for the generation of the chiral ionic liquid's anion, cation, or both.⁵ In our project, the chirality for the ionic liquids was introduced using inexpensive and readily available (L)-lactate and (L)-menthol derivatives. 1-methylpyrrolidine or 1-butylpyrrolidine was heated to reflux in acetonitrile for 48 h with (-)-chloromethyl menthyl ether (Scheme 1).¹⁴ The intermediate chiral ionic salts, 1-butyl-1-menthoxyethylpyrrolidinium chloride and 1-methyl-1-menthoxyethylpyrrolidinium chloride were formed in good yields; 90% and 92% respectively. The spectroscopic analysis of intermediate ionic salts confirmed the purity of the products. The specific rotation was determined by polarimetry. The intermediate

salts were then subjected to anion exchange without further purification. Ionic salts, 1-butyl-1-menthoxyethylpyrrolidinium chloride and 1-methyl-1-menthoxyethylpyrrolidinium chloride were separately reacted with tetrafluoroboric acid, potassium hexafluorophosphate and lithium bis(trifluoromethane)sulfonimide by replacing the chloride anion with tetrafluoroborate, hexafluorophosphate and bis(trifluoromethane)sulfonimide respectively (Scheme 1).¹⁴ These anion exchange reactions resulted six new ionic liquids bearing a chiral cation in good yields (Table 1; **1a–1c** and **1e–1g**). Additionally, the intermediate ionic salts were further reacted with sodium L-lactate by exchanging the chloride anion with L-lactate anion providing two chiral ionic liquids bearing both a chiral anion and a cation (Table 1; **1d** and **1h**). Both 1-butyl-1-menthoxyethylpyrrolidinium L-lactate and 1-methyl-1-menthoxyethylpyrrolidinium L-lactate were semi-solids at room temperature and obtained in good yields. The structure and the purity of final chiral ionic liquids were confirmed by ¹H NMR, ¹³C



Scheme 1. Synthesis of chiral ionic liquids from the reactions of 1-alkylpyrrolidine and (-)-chloromethyl menthyl ether.

Table 1
1-alkyl-1-menthoxyethylpyrrolidinium chiral ionic liquids synthesized in the lab.

Ionic liquid (Yield %)	Ionic liquid (Yield %)
<p>[1a] - R = C₄H₉, (-)-1-butyl-1-menthoxyethylpyrrolidinium tetrafluoroborate (84.5%) [1e] - R = CH₃, (-)-1-methyl-1-menthoxyethylpyrrolidinium tetrafluoroborate (82.3%)</p>	<p>[1b] - R = C₄H₉, (-)-1-butyl-1-menthoxyethylpyrrolidinium hexafluorophosphate (81.8%) [1f] - R = CH₃, (-)-1-methyl-1-menthoxyethylpyrrolidinium hexafluorophosphate (81.0%)</p>
<p>[1c] - R = C₄H₉, (-)-1-butyl-1-menthoxyethylpyrrolidinium bis(trifluoromethane)sulfonimide (89.4%) [1g] - R = CH₃, (-)-1-methyl-1-menthoxyethylpyrrolidinium bis(trifluoromethane)sulfonimide (84.4%)</p>	<p>[1d] - R = C₄H₉, (-)-1-butyl-1-menthoxyethylpyrrolidinium lactate (89.5%) [1h] - R = CH₃, (-)-1-methyl-1-menthoxyethylpyrrolidinium lactate (86.4%)</p>

Download English Version:

<https://daneshyari.com/en/article/5257947>

Download Persian Version:

<https://daneshyari.com/article/5257947>

[Daneshyari.com](https://daneshyari.com)