



# Rhenium-catalyzed deoxydehydration of renewable biomass using sacrificial alcohol as reductant



Justin Gossett, Radhey Srivastava\*

Department of Chemistry, University of Louisiana at Lafayette, Lafayette, LA 70504, USA

## ARTICLE INFO

### Article history:

Received 24 May 2017

Revised 31 July 2017

Accepted 11 August 2017

Available online 23 August 2017

### Keywords:

Deodehydration

Vicinal diol

Reductant


Alkene

Ammonium perrhenate

2,4-dimethyl-3-pentanol

## ABSTRACT

Catalytic deoxydehydration (DODH) of vicinal diols is studied. We find that  $\text{NH}_4\text{ReO}_4$  (ammonium perrhenate, APR) catalyzes the DODH of glycols to alkenes by sacrificial alcohol (2,4-dimethyl-3-pentanol) at 140–165 °C. The product yields range from good to excellent and no isomers detected. The catalytic DODH reaction of glycols to alkene is of potential value for the production of chemicals and fuels from the renewable biomass-derived polyols.



© 2017 Elsevier Ltd. All rights reserved.

## Introduction

The biomass contains abundant hydroxyl groups, which is found in both carbohydrates (obtained from cellulose, hemicellulose, and starch) and glycerol (obtained from triglycerides). The strategy is to reduce oxygen content from biomass to produce diverse classes of value-added chemicals. Initial efforts were focused on dehydration<sup>1</sup> and deoxydehydration (DODH)<sup>2</sup> processes; significant advances accomplished with both, dehydration and DODH processes. To develop a proficient process for the conversion of renewable biomass into reduced oxygen-content products (fuels and value-added chemicals) has fascinated increased attention with the growing demand for sustainability.<sup>3,4</sup> Recent research interest has focused on selective oxygen removal through deoxydehydration (DODH), in which vicinal hydroxyl groups removed to form olefin (Scheme 1). The olefins obtained from DODH reactions are valuable and have diverse applications in the manufacture of chemical intermediates such as polymers and lubricants.


The first report on DODH was disclosed by Andrews and Cook<sup>5</sup> using  $(\text{C}_5\text{Me}_5)\text{ReO}_3$  catalyst in the presence of phosphine

reductants (Scheme 1). A valued contribution discloses by Gable et al.<sup>6</sup> vis-à-vis mechanism of DODH reaction catalyzed by (tris-pyrazolylborate)  $\text{ReO}_3$  in the presence of phosphine as reductant. Lately, significant advances have been achieved using more economical and benign reductant for DODH reaction. The high valent oxorhenium complexes have been used for DODH reaction employing a variety of reductants such as  $\text{H}_2$ <sup>7</sup> and  $\text{Na}_2\text{SO}_3$ .<sup>8</sup> Afterwards various research groups developed Re-catalyzed DODH of vicinal diols<sup>9</sup> in the presence of primary, benzylic, and secondary alcohols as reductants. Sacrificial alcohols have been widely used as effective reductants for the conversion of glycol to olefin in the presence of rhenium-based catalysts.


The Bergman group established a hydrogen-transfer type DODH reaction catalyzed by  $[\text{Re}_2(\text{CO})_{10}]$  and  $[\text{BrRe}(\text{CO})_5]$  in conjunction with a secondary alcohol as solvent/reductant.<sup>10</sup> The Toste group demonstrated the efficient conversion of higher polyols by secondary alcohols to form unsaturated alcohols, unsaturated ethers, and polyenes (as well as the associated ketones) in the presence of  $\text{MeReO}_3$  (MTO) catalyst.<sup>11</sup> Further, Abu Omar and Liu reported the MTO-catalyzed redox disproportionation of glycerol to allyl alcohol, acrolein, and propanal, which can be distilled. In addition to distillable product, they also obtained nonvolatile dihydroxyacetone.<sup>12</sup> Hydroaromatics have also been established as effective reductant for the glycol deoxydehydration in the presence of Re-cat-

\* Corresponding author.

E-mail address: [rss@louisiana.edu](mailto:rss@louisiana.edu) (R. Srivastava).



Scheme 1. Metal-catalyzed DODH reaction.

Scheme 2.  $[\text{Cp}^*\text{Ru}(\text{CO})_2]_2$ -catalyzed hydrodeoxygenation and hydrocracking and  $\text{RuCl}_2(\text{R}_2\text{SO})_4$ -catalyzed hydrogenolysis of diols and epoxides (R = methyl and tetramethylene).

alyst.<sup>13</sup>  $[\text{Cp}^*\text{Ru}(\text{CO})_2]_2$ -catalyzed hydrodeoxygenation and hydrocracking and  $\text{RuCl}_2(\text{R}_2\text{SO})_4$ -catalyzed hydrogenolysis of diols and epoxides has also been established by us (Scheme 2).<sup>14,15</sup>

The Nicholas group<sup>9c</sup> have used ammonium perrhenate (APR) catalyst for the DODH reaction of vicinal diols in the presence of benzylic alcohol as reductant. In the pursuit to develop a new DODH processes for the conversion of renewable cellulosic biomass to valuable chemicals, we revisited the chemistry of  $\text{NH}_4\text{ReO}_4$  and developed a modified process, which requires a small amount of secondary alcohol ( $4.28 \times 10^{-4}$  to  $8.56 \times 10^{-4}$  mol) as a reductant compared to earlier reports<sup>9b,c</sup> (Scheme 3). The reaction is clean and no byproducts except the oxidized byproduct of 2,4-dimethyl-3-pentanol detected.

## Experimental section

### General information

All reagents were obtained commercially and used without further purification. All solvents were ACS grade and were used directly (unless otherwise described in the procedures). GC–MS analyses were performed on an Agilent instrument using a Stabilwax capillary column. NMR spectra were recorded in  $\text{CDCl}_3$  with tetramethylsilane (TMS) as the internal standard for  $^1\text{H}$  (Varian, 400 MHz) and for  $^{13}\text{C}$  (100 MHz) spectra.

### Typical reaction procedure

Glycerol (0.3 mmol, 42 mg),  $\text{NH}_4\text{ReO}_4$  (0.03 mmol, 8 mg), 2,4-dimethyl-3-pentanol (0.43–0.86 mmol, 60–120  $\mu\text{L}$ ), and 1 mL anhydrous toluene were added to a thick-walled Ace glass reactor tube. The Teflon seal was closed and the reactor was placed in a heating mantle connected with a Digi Troll (Glas-Col) digital temperature controller equipped with a thermocouple at 140–165  $^\circ\text{C}$  for 5–48 h while stirring. After cooling to room temperature, the solution was filtered to remove the precipitated  $\text{NH}_4\text{ReO}_4$  and analyzed with GC–MS.



Scheme 3. Optimized DODH reaction scheme for glycol conversion to the corresponding olefins.

Table 1

Optimization of solvents and alcohol reductants in the presence of  $\text{NH}_4\text{ReO}_4$  catalysts.

Entry	Solvent	Styrene (%)	Time (h)
1	Toluene	84	4
2	Toluene	99	5
3	1-Butanol	42	24
4	Benzene	17	24
5	THF	32	24
6	Acetonitrile	19	24
7	3-Octanol	75	24
8	Isopropanol	10	24

### Isolation

$\text{NH}_4\text{ReO}_4$  (28 mg, 0.1 mmol), (+)-diethyl tartrate (198 mg, 1 mmol), 2,4-dimethyl-3-pentanol (388  $\mu\text{L}$ ), and anhydrous toluene (6 mL) were added to a thick-walled Ace glass reactor tube. The reactor was placed in a heating mantle bath in the range of 165  $^\circ\text{C}$  for 24 h while stirring magnetically. The reaction mixture was cooled and filtered over silica gel. Solvents were removed using a rotary evaporator under reduced pressure. The product was separated by column chromatography using ethyl acetate/hexane eluent. Product yield was 73 mg, 42%.

## Result and discussion

In pursuit to improve catalytic systems for DODH with better efficiency, we prefer to use ionic ammonium perrhenate (APR,  $\text{NH}_4\text{ReO}_4$ ) as a suitable catalyst. APR is inexpensive, known for DODH ability, greater hydrolytic stability compared to  $\text{MeReO}_3$ ,<sup>16</sup> and its ionic nature. To test the viability of alcohols as a reductant for the DODH of glycols, an exploratory reaction was carried out with styrene diol, 2,4-dimethyl-3-pentanol and ammonium perrhenate (10 mol%) in anhydrous toluene, which produced styrene quantitatively in 5 h at 165  $^\circ\text{C}$ . We began the survey with polar and apolar solvents at various temperatures. The reaction performed in anhydrous toluene produced an excellent yield of the alkene whereas hydrated toluene and benzene resulted in a low yield. The addition of 4 Å molecular sieve did not change the product yield. The coordinating solvent such as THF and acetonitrile also produced significantly lower yield of styrene, probably due to the coordination of solvent to Re (Table 1). Our next step was to screen the effect of temperature and we found that 160–165  $^\circ\text{C}$  was the optimum temperature for the model substrate for optimum yield of alkene. We preferred secondary alcohol (2,4-dimethyl-3-pentanol) over butanol-1,3-octanol, and isopropanol for DODH reductant because the higher yield of alkenes and the resulting ketone co-product could be easily separated from the olefinic products in GC–MS.

After optimizing reaction conditions (10 mol% APR, 0.3 mmol glycol, 2,4-dimethyl-3-pentanol in anhydrous toluene, 140–165  $^\circ\text{C}$ , 4–24 h), we determined that the representative glycols were converted to the corresponding olefins in good to excellent yields (Scheme 3, Table 2). All polyols subjected to catalytic reaction are efficiently converted to the corresponding alkenes regioselectively as is shown in Table 2. The activated diol such as 1-phenyl-1, 2-ethanediol converted to styrene quantitatively in 5 h at 165  $^\circ\text{C}$  (run 1, Table 2). Long chain diols, tetradecanediol

Download English Version:

<https://daneshyari.com/en/article/5258913>

Download Persian Version:

<https://daneshyari.com/article/5258913>

[Daneshyari.com](https://daneshyari.com)