



Rapid cycloaddition of a diazo group with an unstrained dipolarophile



Matthew R. Aronoff^{a,†}, Brian Gold^{a,†}, Ronald T. Raines^{a,b,*}

^a Department of Chemistry, University of Wisconsin–Madison, Madison, WI 53706, USA

^b Department of Biochemistry, University of Wisconsin–Madison, Madison, WI 53706, USA

ARTICLE INFO

Article history:

Received 8 March 2016

Revised 3 April 2016

Accepted 7 April 2016

Available online 28 April 2016

Keywords:

Diazo group

1,3-Dipolar cycloaddition

(*E*)-Ethyl 4,4,4-trifluorobut-2-enoate

Hydrogen bond

Hyperconjugation

ABSTRACT

The cycloaddition of a diazoacetamide with ethyl 4,4,4-trifluorocrotonate proceeds with $k = 0.1 \text{ M}^{-1} \text{ s}^{-1}$. This second-order rate constant rivals those of optimized strain-promoted azide–alkyne cycloadditions, even though the reaction does not release strain. The regioselectivity and a computational distortion/interaction analysis of the reaction energetics are consistent with the formation of an N–H...F–C hydrogen bond in the transition state and the electronic character of the trifluorocrotonate. Analogous reactions with an azidoacetamide dipole or with an acrylate or crotonate dipolarophile are much slower. These findings suggest a new strategy for the design of diazo-selective reagents for chemical biology.

© 2016 Elsevier Ltd. All rights reserved.

Introduction

The diazo group has the potential to be a valuable addition to the armamentarium of chemical biology. A complement to azido groups, diazo groups have been employed in many biochemical applications, including reversible¹ and non-reversible protein alkylations,² nucleic acid conjugation,³ and other bioconjugations.⁴ Extant strategies to install pendant diazo groups⁵ and to prepare them directly by deimidogenation of a parent azide⁶ facilitate their use.

Mistakenly believed to be too reactive for physiological investigations, a diazo moiety stabilized by electronic delocalization can tolerate metabolism while retaining reactivity.⁷ Nevertheless, the use and development of diazo compounds in chemical biology are constrained by a paucity of chemoselective reactions. Most confounding is the overlapping reactivity of the diazo group with that of the azido group in strain-promoted azide–alkyne cycloadditions (SPAACs).^{7,8}

The need for reactions that are chemoselective in a biological context continues to grow.⁹ Recently, we explored 1,3-dipolar cycloadditions of diazo compounds in the presence of azides.¹⁰ Although we were able to achieve diazo group-specific conjugation in the absence of a catalyst (Scheme 1), the reaction rates were

low—comparable to those obtained with early SPAAC reagents and exceeded by new SPAAC reagents.^{8c,11} Hence, we sought a faster cycloaddition.

We were aware that fluoro groups can elicit favorable orbital interactions in cycloadditions¹² and other reactions.¹³ Accordingly, we explored the utility of fluoro groups in a cycloaddition reaction with a diazo compound. We discovered that the reaction of a diazoacetamide with a fluorinated crotonate proceeds at an extraordinary rate. We used computations to provide insight into this rate acceleration. The results reveal a general strategy for the design of efficacious diazo-selective reagents for applications in chemical biology.


Results and discussion

In an initial screen of reactions, we dissolved *N*-benzyl-2-diazoacetamide (**1**) in 1:1 CH₃CN/H₂O and added ethyl 4,4,4-trifluorocrotonate (**2**) to the resulting solution (Scheme 2).¹⁴ We found that the diazo compound was consumed rapidly, as monitored with thin-layer chromatography. The reaction product had the mass of the pyrazoline that would result from the dipolar cycloaddition reaction of the diazo group and the alkene. We monitored the reaction by ultraviolet spectroscopy, and found the observed rate constant to be $k = 0.105 \text{ M}^{-1} \text{ s}^{-1}$. This rate constant is comparable to that of an azide with BARAC, which is the fastest known SPAAC reaction.¹⁵

* Corresponding author. Tel.: +1 608 262 8588; fax: +1 608 890 2583.

E-mail address: rtraines@wisc.edu (R.T. Raines).

† These authors contributed equally to this work.



Scheme 1. Chemoselective cycloaddition of a diazoacetamide with an acrylate ester.


The putative pyrazoline product contained a mixture of isomers. We were able to purify the major product (**3**) and confirm its structure using 3J coupling values from ^1H NMR spectroscopy as verified with heteronuclear multiple bond correlation (HMBC). Based on isolated yields, the cycloaddition occurred to produce **3** and **4** in a 9:1 ratio. This strong regiochemical preference for the *anti* orientation of the carbonyl groups is consistent with the π -acceptor ester group being a stronger electron-withdrawing substituent than is the σ -acceptor trifluoromethyl group and with the formation of an $\text{N}-\text{H}\cdots\text{F}-\text{C}$ hydrogen bond, as has been observed in other contexts.¹⁶

To verify the identity of products **3** and **4**, we sought to oxidize the enantiomeric mixture of pyrazolines to their aromatic pyrazole

equivalents.¹⁷ Attempts at oxidation with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) were unsuccessful, but treatment with Br_2 produced the pyrazole product (Scheme 2). To confirm the assignment of pyrazolines **3** and **4**, the oxidation products were compared to those resulting directly from the reaction of diazoacetamide **1** with ethyl 4,4,4-trifluorobut-2-ynoate (**5**). The ^1H NMR and ^{13}C NMR spectra of **6** and **7** produced by the two routes were indistinguishable.

Next, we confirmed the importance of electronic acceleration provided by allylic fluorination by assessing the cycloaddition of diazoacetamide **1** with ethyl acrylate (**8**), which lacks the trifluoromethyl group, and methyl crotonate (**9**), which lacks the three fluoro groups. Previously, we reported that the reaction of **1** and **8** proceeds with a rate constant of $k = 1.6 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$ (Fig. 1), which is 10^2 -fold less than that of trifluorocrotonate **2**.¹⁰ Here, we found the rate constant for cycloaddition with crotonate **9** to be $k = 5.8 \times 10^{-7} \text{ M}^{-1} \text{ s}^{-1}$, which is 10^5 -fold less than that for trifluorocrotonate **2**. Notably, the effect of allylic fluorination in this reaction is much larger than is the effect of propargylic fluorination on the cycloaddition of azides (~ 400 -fold).^{12d} Finally, when added to an equimolar mixture of diazoacetamide **1** and *N*-benzyl-2-azidoacetamide (**10**), excess trifluorocrotonate **2** reacted only with diazoacetamide **1**, demonstrating chemoselectivity.

We employed computational methods to investigate the origin of the increased reactivity of trifluorocrotonate **2**. Optimizations were performed with Gaussian 09 software¹⁸ at the M06-2X level of theory¹⁹ and the 6-31+G(2d,p) basis set. M06-2X has been shown to describe trends in reactivity accurately in other cycloadditions.²⁰ Optimizations were performed in the gas phase, and



Scheme 2. Rapid cycloaddition of a diazoacetamide with a 4,4,4-trifluorocrotonate ester, and product analysis.




Figure 1. Experimental rate constants and computational parameters of the transition state for cycloadditions of a diazoacetamide with a trifluorocrotonate (**1-2 TS**), acrylate (**1-8 TS**), and unfluorinated crotonate (**1-9 TS**). Optimized geometries, activation energies, distortion/interaction energies, and NBO charges on the dipolarophile (*italics*) were calculated at the M06-2X/6-31+G(2d,p) level of theory. Energies (kcal/mol) include solvation corrections (water) on gas-phase geometries using IEFPCM model (radii = UFF). The methyl ester of trifluorocrotonate **2** was used to model the ethyl ester. Rate constants were measured in 1:1 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (**1 + 2**; **1 + 8**) or CH_3CN (**1 + 9**). Data for the cycloaddition of **1** and **8** were reported previously.¹⁰

Download English Version:

<https://daneshyari.com/en/article/5258948>

Download Persian Version:

<https://daneshyari.com/article/5258948>

[Daneshyari.com](https://daneshyari.com)